

FINITE COMPLEX REFLECTION ARRANGEMENTS ARE $K(\pi, 1)$

DAVID BESSIS

ABSTRACT. Let V be a finite dimensional complex vector space and $W \subseteq \mathrm{GL}(V)$ be a finite complex reflection group. Let V^{reg} be the complement in V of the reflecting hyperplanes. We prove that V^{reg} is a $K(\pi, 1)$ space. This was predicted by a classical conjecture, originally stated by Brieskorn for complexified real reflection groups. The complexified real case follows from a theorem of Deligne and, after contributions by Nakamura and Orlik-Solomon, only six exceptional cases remained open. In addition to solving this six cases, our approach is applicable to most previously known cases, including complexified real groups for which we obtain a new proof, based on new geometric objects. We also address a number of questions about $\pi_1(W \backslash V^{\mathrm{reg}})$, the braid group of W . This includes a description of periodic elements in terms of a braid analog of Springer's theory of regular elements.

CONTENTS

Introduction	2
Notations	6
Topological conventions	6
1. Complex reflection groups, discriminants, braid groups	7
2. Well-generated complex reflection groups	9
3. Symmetric groups, configurations spaces and classical braid groups	12
4. Affine Van Kampen method	14
5. Lyashko-Looijenga coverings	15
6. Tunnels, labels and Hurwitz rule	18
7. Reduced decompositions of Coxeter elements	23
8. The dual braid monoid	27
9. Chains of simples	30
10. The universal cover of $W \backslash V^{\mathrm{reg}}$	34
11. Centralisers of regular elements	38
12. Periodic elements in $B(W)$	49
13. Generalised non-crossing partitions	51
Acknowledgements	53
References	53

INTRODUCTION

Let V be a finite dimensional complex vector space and $W \subseteq \mathrm{GL}(V)$ be a complex reflection group (all reflection groups considered here are assumed to be finite).

Let V^{reg} be the complement in V of the reflecting hyperplanes. In the case when W is a type A reflection group, Fadell and Neuwirth proved in the early 1960's that V^{reg} is a $K(\pi, 1)$ (this is an elementary use of fibration exact sequences, see [28]). Brieskorn conjectured in 1971, [16], that the $K(\pi, 1)$ property holds when W is a complexified real reflection group. It is not clear who first stated the conjecture in the context of arbitrary complex reflection groups. It may be found, for example, in Orlik-Terao's book:

Conjecture 0.1 ([36], p. 163 & p. 259). *The universal cover of V^{reg} is contractible.*

Our main result is a proof of this conjecture. It is clearly sufficient to consider the case when W is irreducible, which we assume from now on. Irreducible complex reflection groups have been classified by Shephard-Todd, [42].

The complexified real case (*i.e.*, Brieskorn's conjecture) was quickly settled by Deligne, [25]. The rank 2 complex case is trivial. The case of the infinite family $G(de, e, n)$ was solved in 1983 by Nakamura, [33] (here again, the monomiality of the group allows an efficient use of fibrations). A few other cases immediately follow from the observation by Orlik-Solomon, [35], that certain discriminants of non-real complex reflection groups are isomorphic to discriminants of complexified real reflection groups.

Combining all previously known results, the conjecture remained open for six exceptional types: G_{24} , G_{27} , G_{29} , G_{31} , G_{33} and G_{34} . We complete the proof of the conjecture by dealing with these cases.

Let $d_1 \leq \dots \leq d_n$ be the degrees of W . Let $d_1^* \geq \dots \geq d_n^* = 0$ be the codegrees of W . We say that W is a *duality* group if $d_i + d_i^* = d_n$ for all i (by analogy with the real case, we then say that d_n is the *Coxeter number* of W , denoted by h). We say that W is *well-generated* if it may be generated by n reflections. Orlik-Solomon observed, by inspecting the classification of Shephard-Todd, that

$$W \text{ is a duality group} \Leftrightarrow W \text{ is well-generated.}$$

The first ten sections of this article are devoted to the proof of the following theorem:

Theorem 0.2. *Let W be a well-generated complex reflection group. The universal cover of V^{reg} is contractible.*

The proof relies on combinatorial and geometric objects which are specific to well-generated groups. It is essentially “case-free”, although a few combinatorial lemmas still require some limited use of the Shephard-Todd classification.

Five of the six open cases are well-generated: G_{24} , G_{27} , G_{29} , G_{33} and G_{34} . The theorem also applies to the complexified real case, for which we obtain a new proof, not relying on [25].

The remaining case, G_{31} , is not well-generated: it is an irreducible complex reflection group of rank 4 which cannot be generated by less than 5 reflections. Fortunately, we may view it as the centraliser of a 4-regular element (in the sense of Springer, [41]) in the group G_{37} (the complexification of the real group of type E_8). By refining the geometric and combinatorial tools introduced in the study of the duality case, one obtains a relative version of Theorem 0.2:

Theorem 0.3. *Let W be a well-generated complex reflection group. Let d be a Springer regular number, let ζ be a primitive complex d -th root of unity, let $w \in W$ be a ζ -regular element. Let $V' := \ker(w - \zeta)$, let W' be the centraliser of w in W , viewed as a complex reflection group acting on V' (see [41]). Let V'^{reg} be the associated hyperplane complement. The universal cover of V'^{reg} is contractible.*

This in particular applies to G_{31} and, based on earlier results, completes the proof of the $K(\pi, 1)$ conjecture.

Note that the case $d = 1$ in Theorem 0.3 is precisely Theorem 0.2. We state the two results separately because it better reflects the organisation of the proof.

As by-products of our construction, we obtain new cases of several standard conjectures about the braid group of W , defined by

$$B(W) := \pi_1(W \setminus V^{\text{reg}}).$$

Proving Theorem 0.2: general strategy. The general architecture of our proof is borrowed from Deligne’s original approach but the details are quite different. Every construction here is an analogue of a construction from [25] but relies on different combinatorial and geometric objects. Like in [25], one studies a certain braid monoid M , whose structure expresses properties of reduced decompositions in W , and one proves that it is a lattice for the divisibility order (this amounts to saying that the monoid is *Garside*). Like in [25], one uses semi-algebraic geometry to construct an open covering of the universal cover of V^{reg} , with the property that non-empty intersections are contractible. This implies that the universal cover is homotopy equivalent to the nerve of the covering. Like in [25], one interprets this nerve as a certain flag complex obtained from M . Like in [25], the contractibility of the nerve follows from the lattice property for M . However, our proof does not use the classical braid monoid, but a dual braid monoid ([3], [6]), whose construction is generalised to all well-generated complex reflection groups. The construction of the open covering is the most problematic step: by contrast with the real case, one cannot rely on the notions of *walls* and *chambers*. The idea here is to work in $W \setminus V^{\text{reg}}$ and to use a generalisation of the Lyashko-Looijenga morphism. This morphism allows a description of $W \setminus V^{\text{reg}}$ by means of a ramified covering of a type A reflection orbifold. Classical objects like *walls*, *chambers* and *galleries* can somehow be “pulled-back”, via the Lyashko-Looijenga morphism, to give semi-algebraic objects related to the dual braid monoid.

Lyashko-Looijenga coverings. The quotient map $\pi : V^{\text{reg}} \rightarrow W \setminus V^{\text{reg}}$ is a regular covering. Once a system of basic invariants $(f_1, \dots, f_n)$ is chosen, the quotient space $W \setminus V^{\text{reg}}$ identifies with the complement in $\mathbb{C}^n$ of an algebraic hypersurface $\mathcal{H}$, the *discriminant*, of equation $\Delta \in \mathbb{C}[X_1, \dots, X_n]$. If W is an irreducible duality complex reflection group, it is possible to choose $(f_1, \dots, f_n)$ such that

$$\Delta = X_n^n + \alpha_2 X_n^{n-2} + \dots + \alpha_n,$$

where $\alpha_2, \dots, \alpha_n \in \mathbb{C}[X_1, \dots, X_{n-1}]$. Let $Y := \text{Spec } \mathbb{C}[X_1, \dots, X_{n-1}]$, together with the natural map $p : W \setminus V^{\text{reg}} \rightarrow Y$. We have an identification $W \setminus V \simeq \mathbb{C}^n \xrightarrow{\sim} \mathbb{C} \times Y$ sending the orbit $\bar{v}$ of $v \in V$ to $(f_n(v), p(\bar{v}))$. The fiber of p over $y \in Y$ is a line L_y which intersects $\mathcal{H}$ at n points (counted with multiplicities). Generically, the n points are distinct. Let $\mathcal{K}$ be the *bifurcation locus*, i.e., the algebraic hypersurface of Y consisting of points y such that the intersection has cardinality $< n$. Classical results from invariant theory of complex reflection groups make possible (and very easy) to generalise a construction by Looijenga and Lyashko: the map LL (for “Lyashko-Looijenga”) sending $y \in Y - \mathcal{K}$ to the subset $\{x_1, \dots, x_n\} \subseteq \mathbb{C}$ such that $p^{-1}(y) \cap \mathcal{H} = \{(x_1, y), \dots, (x_n, y)\}$ is a regular covering of degree $n!h^n/|W|$ of the (centered) configuration space of n points in $\mathbb{C}$. In particular, $Y - \mathcal{K}$ is a $K(\pi, 1)$. This observation, which is apparently new in the non-real case, already allows a refinement of our earlier results ([2], [8]) on presentations for the braid group of W .

The dual braid monoid. When W is complexified real, a *dual braid monoid* was constructed in [3] (generalising the construction of Birman-Ko-Lee, [11]; similar partial results were independently obtained by Brady-Watt, [13]). The construction was later generalised in [6] to the

complex reflection group $G(e, e, n)$. Let R be the set of all reflections in a well-generated group W . The idea is that the pair (W, R) has some “Coxeter-like” features. Instead of looking at relations of the type

$$\underbrace{sts\dots}_{m_{s,t}} = \underbrace{tst\dots}_{m_{s,t}},$$

one considers relations of the type

$$st = tu$$

where $s, t, u \in R$. Let $\mathcal{S}$ be the set of all relations of this type holding in W . In general, $B(W) \not\simeq \langle R | \mathcal{S} \rangle$, but it is possible to find natural subsets $R_c \subseteq R$ and $\mathcal{S}_c \subseteq \mathcal{S}$ such that $B(W) \simeq \langle R_c | \mathcal{S}_c \rangle$ (if W is complexified real, $R_c = R$). The elements of $\mathcal{S}_c$ are called *dual braid relations*. The choices of R_c and $\mathcal{S}_c$ are natural once a *Coxeter element* c has been chosen (the notion of *Coxeter element* generalises to the non-real well-generated groups in terms of Springer’s theory of regular elements). Since the relations are positive, one may view the presentation as a monoid presentation, defining a monoid $M(W)$. The crucial property of this monoid is that it is a lattice for the divisibility order or, more precisely, a *Garside monoid*. Following Deligne, Bestvina, T. Brady and Charney-Meier-Whittlesey, the Garside structure provides a convenient simplicial Eilenberg-McLane $K(B(W), 1)$ space ([25], [10], [12], [21]). The earlier results on the dual braid monoid are improved here in two directions:

- The construction is generalised to the few exceptional cases (G_{24} , G_{27} , G_{29} , G_{33} and G_{34}) not covered by [3] and [6].
- A new geometric interpretation is given, via the Lyashko-Looijenga covering. This interpretation is different from the one given in [3, Section 4].

The second improvement is the most important. It relies (so far) on a counting argument, following and extending a property which, for the complexified real case, was conjectured by Looijenga and proved in a letter from Deligne to Looijenga, [26].

Tunnels. The classical theory of real reflection groups combines a “combinatorial” theory (Coxeter systems) and a “geometric” theory (expressed in the language, invented by Tits, of walls, chambers, galleries, buildings...) We expect the dual braid monoid approach to eventually provide effective substitutes for much of this classical theory. A first step in this direction is the notion of *tunnel*, which is a rudimentary geometric object replacing the classical notion of *minimal gallery between two chambers*. An important difference with the classical geometric language is that tunnels are naturally visualised in $W \backslash V$ (instead of V). A tunnel T is a path in $W \backslash V^{\text{reg}}$ drawn inside a single line L_y (for some $y \in Y$) and with constant imaginary part. It represents an element b_T of the dual braid monoid M . An element of M is *simple* if it is represented by a tunnel. This notion coincides with the notion of simple element associated with the Garside structure. In the classical approach, for any chamber $\mathcal{C}$, there are as many equivalence classes of minimal galleries starting at $\mathcal{C}$ as simple elements (this number is $|W|$). Here the situation is different: in a given L_y , not all simples are represented. The simples represented in different L_y ’s may be compared thanks to a huge “basepoint” $\mathcal{U}$ which is both dense in $W \backslash V^{\text{reg}}$ and contractible.

Proving Theorem 0.3. The strategy is the same as for Theorem 0.2. With the notations of the theorem, the quotient space $W' \backslash V'^{\text{reg}}$ may be identified with $(W \backslash V^{\text{reg}})^{\mu_d}$, for the natural action of the cyclic group μ_d . This action induces an automorphism of $B(W)$ which, unfortunately, does not preserve the dual braid monoid. However, it is possible to replace $B(W)$ by a sort of categorical barycentric subdivision, its *d-divided Garside category* M_d , on which μ_d acts *diagram automorphisms*. This construction is explained in my separate article [5]. The fixed subcategory $M_d^{\mu_d}$ is again a Garside category. It should be thought of a *dual braid category* for

$B(W')$ and gives rise to a natural simplicial space, whose realisation is an Eilenberg-MacLane space. As before, one shows that $(W \setminus V^{\text{reg}})^{\mu_d}$ is homotopy equivalent to this simplicial model, by studying the nerve of a certain open covering of a certain model of the universal cover of $(W \setminus V^{\text{reg}})^{\mu_d}$, very similar to the one used for $W \setminus V^{\text{reg}}$ (except that one has to replace the contractible “basepoint” $\mathcal{U}$ by a family of non-overlapping contractible “basepoints”, one for each object of $M_d^{\mu_d}$.) This involves replacing tunnels by a suitable notion of *circular tunnels*. This article focuses the geometric aspects of the proof of Theorem 0.3 – it is probably fair to say that the true explanation lies in the properties of M_d and in the general theorems about periodic elements in Garside groupoids that are explained in [5].

By-products.

Theorem 0.4. *Braid groups of well-generated complex reflection groups are Garside groups.*

In the situation of Theorem 0.3, we prove that the braid group $B(W')$ is a weak Garside group, which is almost as good.

In particular, $B(W)$ is torsion-free, admits nice solutions to the word and conjugacy problems, is biautomatic, admits a finite $K(\pi, 1)$ (our construction provides an explicit one), and much more – see [23] for a quite complete reference. None of this was known for the six exceptional groups mentioned above.

Theorem 0.5. *The centre of the braid group of an irreducible well-generated complex reflection group is cyclic.*

Again, the cases of G_{24} , G_{27} , G_{29} , G_{33} and G_{34} are new. It was conjectured by Broué-Malle-Rouquier, [18], and should also hold for non-well-generated groups. The only case left, G_{31} , is probably not too difficult to solve from our description.

For $B(G_{29})$, $B(G_{31})$, $B(G_{33})$ and $B(G_{34})$, no presentations were known until now, although some conjectures made in [8] were supported by strong evidences.

Theorem 0.6. *The conjectural presentations for $B(G_{29})$, $B(G_{31})$, $B(G_{33})$ and $B(G_{34})$ given in [8] are correct.*

Combined with [7] and [8], this completes the longstanding task of finding presentations for all generalised braid groups associated with finite complex reflection groups. Theorem 0.6 is much easier than the previously mentioned results and only relies on a minor improvement over [2] and [8]. However, the material presented here allows for a more conceptual proof.

Periodic elements in braid groups. In connection with their work on Deligne-Lusztig varieties (see [17] for more details), Broué-Michel predicted the existence of an analog for braid groups of Springer’s theory of regular elements. This amounts to a conjectural description of periodic elements (elements with a central power) and their centralisers. When W is the symmetric group, periodic elements in $B(W)$ may be understood thanks to Kerékjártó’s theorem on periodic homeomorphisms of the disk. In the more general setting of spherical type Artin groups, finding a simple description of periodic elements was an open question. We are able to solve these problems when W is well-generated: Theorem 12.5 contains a complete description of the roots of the generator of the center of the pure braid group $P(W)$ and of their centralisers.

As for Theorem 0.3, the main conceptual ingredient towards the proof of Theorem 12.5 is a general property of Garside categories, explained in our separate paper [5]. What is done here is the minor step consisting of re-interpreting the general *Kerékjártó theorem for Garside categories* from [5] in terms of the S^1 -structure on the regular orbit space $W \setminus V^{\text{reg}}$.

Non-crossing partitions. A combinatorial by-product of our approach is a general construction of *generalised non-crossing partitions*, associated to each type of well-generated complex reflection groups.

In the classical cases A_n , B_n , D_n and, more generally, $G(e, e, n)$, the structure of $M(W)$ is understood in terms of suitable notions of *non-crossing partitions* ([1], [6], [7]). The dual braid monoid of an irreducible well-generated complex reflection W gives rise to a lattice of *generalised non-crossing partitions*, whose cardinal is the *generalised Catalan number*

$$\text{Cat}(W) := \prod_{i=1}^n \frac{d_i + d_n}{d_i}.$$

(the term “partition” should not be taken too seriously: except for the classical types, lattice elements do not have natural interpretations as actual set-theoretic partitions.) It is likely that this combinatorial object has some representation-theoretic interpretation. In the “badly-generated” case, $\text{Cat}(W)$ may fail to be an integer, and the natural substitute for $NCP(W)$ is the graph of simples of the dual braid category.

NOTATIONS

We save the letter i for indexing purposes and denote by $\sqrt{-1}$ a complex square root of -1 fixed once for all. If n is a positive integer, we denote by ζ_n the standard n -th root of unity $\exp(2\sqrt{-1}\pi/n)$.

Many objects depend on a complex reflection group W , e.g., the braid group $B(W)$. We often drop the explicit mention of W , and write B for $B(W)$. When n is an integer, we denote by B_n the braid group on n strings, together with its standard generating set $\sigma_1, \dots, \sigma_{n-1}$; it is isomorphic to the braid group of $\mathfrak{S}_n$ in its permutation reflection representation (see Section 3). The groups B and B_n appear simultaneously and should not be confused.

TOPOLOGICAL CONVENTIONS

Let E be a topological space. Let γ be a path in E , *i.e.*, a continuous map $[0, 1] \rightarrow E$. We say that γ is a *path from* $\gamma(0)$ *to* $\gamma(1)$ – or that $\gamma(0)$ is the *source* and $\gamma(1)$ the *target*. The concatenation rule is as follows: if γ, γ' are paths such that $\gamma(1) = \gamma'(0)$, the product $\gamma\gamma'$ is the path mapping $t \leq 1/2$ to $\gamma(2t)$ and $t \geq 1/2$ to $\gamma'(2t - 1)$.

A path γ from e to e' induces an isomorphism $\phi_\gamma : \pi_1(E, e) \xrightarrow{\sim} \pi_1(E, e')$.

Let X be a subspace of E . Assume that X is simply-connected (in all applications, X will actually be contractible). Then for all $x, x' \in X$ there is a unique homotopy class of paths from x to x' drawn in X , thus a natural isomorphism $\phi_{X, x, x'} : \pi_1(E, x) \xrightarrow{\sim} \pi_1(E, x')$. By unicity, $\phi_X := (\phi_{X, x, x'})_{x, x' \in X}$ is a transitive system of isomorphisms between the $(\pi_1(E, x))_{x \in X}$. We denote by

$$\pi_1(E, X)$$

the transitive limit. Practically speaking, $\pi_1(E, X)$ should be thought of as any $\pi_1(E, x)$, for some $x \in X$, together with an unambiguous recipe thanks to which any path in E from any $x' \in X$ to any $x'' \in X$ represents a unique element of $\pi_1(E, x)$ – moreover, one may forget about which $x \in X$ was chosen and change it at our convenience.

Assume that E is locally simply-connected. To get a model $\tilde{E}_x$ of the universal cover $\tilde{E}$, one chooses a basepoint $x \in E$. A path γ from x to $e \in E$ represents a point $\bar{\gamma} \in \tilde{E}_x$ in the fiber over e . Two paths represent the same point if and only if they are homotopic (with fixed endpoints). The topology of $\tilde{E}_x$ around $\bar{\gamma}$ is described by concatenation at the target with paths starting at e and drawn in simply-connected neighbourhoods of e . The fundamental group $\pi_1(E, x)$ acts

on the left on $\tilde{E}_x$ by concatenation at the source. More generally, a path from x to x' induces an isomorphism $\tilde{E}_x \xrightarrow{\sim} \tilde{E}_{x'}$. Here again, if X is simply-connected, we have a transitive system of isomorphisms between $(\tilde{E}_x)_{x \in X}$ whose limit is a model $\tilde{E}_X$ of the universal cover. Practically speaking, any path from any $x \in X$ to any $e \in E$ unambiguously represents a point in $\tilde{E}_X$ (in the fiber over e). Moreover, $\pi_1(E, X)$ acts on the left on $\tilde{E}_X$.

1. COMPLEX REFLECTION GROUPS, DISCRIMINANTS, BRAID GROUPS

Let V be a vector space of finite dimension n . A *reflection group* in $\mathrm{GL}(V)$ is a subgroup W generated by (*generalised*) *reflections*, i.e., elements whose fixed subspace is a hyperplane. When the basefield of V is $\mathbb{C}$, we say that W is a *complex reflection group*. We are only interested in finite reflection groups and will always assume finiteness, unless otherwise specified.

Let $W \subseteq \mathrm{GL}(V)$ be a complex reflection group. A *system of basic invariants* for W is an n -tuple $f = (f_1, \dots, f_n)$ of homogeneous generators of $\mathcal{O}_V^W$, the algebra of W -invariant polynomial functions on V . A classical theorem of Shephard-Todd, [42], asserts that such tuples exist, and that they consist of algebraically independent terms. Set $d_i := \deg f_i$; these numbers are the *degrees* of W . Up to reordering, we may assume that $d_1 \leq d_2 \leq \dots \leq d_n$. The sequence $(d_1, \dots, d_n)$ is then independent of the choice of f .

Choosing a system of basic invariants f amounts to choosing a graded algebra isomorphism $\mathcal{O}_V^W \simeq \mathbb{C}[X_1, \dots, X_n]$, $f_i \mapsto X_i$, where the indeterminate X_i is declared homogeneous with degree d_i . Geometrically, this isomorphism identifies the categorical quotient $W \backslash V$ with the affine space $\mathbb{C}^n$.

Further features of the invariant theory of complex reflection groups involve invariant vector fields and invariant differential forms on V .

Theorem 1.1 ([36], Lemma 6.48). *The $\mathcal{O}_V^W$ -modules $(\mathcal{O}_V \otimes V)^W$ and $(\mathcal{O}_V \otimes V^*)^W$ are free of rank n .*

If $f = (f_1, \dots, f_n)$ is a system of basic invariants, $df := (df_1, \dots, df_n)$ is a $\mathcal{O}_V^W$ -basis for $(\mathcal{O}_V \otimes V^*)^W$. Being homogeneous, the module $(\mathcal{O}_V \otimes V)^W$ admits an homogeneous basis.

Definition 1.2. A system of basic derivations for W is an homogeneous $\mathcal{O}_V^W$ -basis $\xi = (\xi_1, \dots, \xi_n)$ of $(\mathcal{O}_V \otimes V)^W$, with $\deg(\xi_1) \geq \deg(\xi_2) \geq \dots \geq \deg(\xi_n)$.

The sequence $(d_1^*, \dots, d_n^*) := (\deg(\xi_1), \dots, \deg(\xi_n))$ is the sequence of *codegrees* of W (it does not depend on the choice of ξ).

Note that, as in [2], we label codegrees in decreasing order, which is slightly unusual. When W is a complexified real reflection group, we have $V \simeq V^*$ as W -modules, thus $d_i^* = d_{n-i+1} - 2$ for all i . This relation is specific to the real situation and is not relevant here.

The Euler vector field on V is invariant and of degree 0. Thus $d_n^* = 0$.

Invariant vector fields define vector fields on the quotient variety. Let f be a system of basic invariants and ξ be a system of basic derivations. For $j \in \{1, \dots, n\}$, the vector field ξ_j defines a vector field $\bar{\xi}_j$ on $W \backslash V$. Since $\frac{\partial}{\partial f_1}, \dots, \frac{\partial}{\partial f_n}$ is a $\mathcal{O}_V^W$ -basis of the module of polynomial vector fields on $W \backslash V$, we have

$$\bar{\xi}_j = \sum_{i=1}^n m_{i,j} \frac{\partial}{\partial f_i},$$

where the $m_{i,j}$ are uniquely defined elements of $\mathcal{O}_V^W$.

Definition 1.3. The discriminant matrix of W (with respect to f and ξ) is $M := (m_{i,j})_{i,j}$.

By weighted homogeneity, one has:

Lemma 1.4. *For all i, j , $\text{wt}(m_{i,j}) = d_i + d_j^*$.*

The vector space V decomposes as a direct sum $\bigoplus_i V_i$ of irreducible representations of W . Denote by W_i the irreducible reflection group in $\text{GL}(V_i)$ generated by (the restriction of) the reflections in W whose hyperplanes contain $\bigoplus_{j \neq i} V_j$. We have $W \simeq \prod_i W_i$. Viewing W and the W_i 's as *reflection groups*, i.e., groups endowed with a reflection representation, it is natural to actually write $W = \bigoplus_i W_i$.

We denote by $\mathcal{A}$ the arrangement of W , i.e., the set of reflecting hyperplanes of reflections in W . We set

$$V^{\text{reg}} := V - \bigcup_{H \in \mathcal{A}} H.$$

Denote by p the quotient map $V \rightarrow W \backslash V$. Choose a basepoint $v_0 \in V^{\text{reg}}$.

Definition 1.5 ([18]). The *braid group* of W is $B(W) := \pi_1(W \backslash V^{\text{reg}}, p(v_0))$.

Later on, when working with well-generated reflection groups, we will slightly upgrade this definition, by replacing the basepoint by a convenient contractible subspace of $W \backslash V^{\text{reg}}$ (see Definition 6.3).

To write explicit equations, one chooses a system of basic invariants f . The *discriminant* $\Delta(W, f) \in \mathbb{C}[X_1, \dots, X_n]$ is the reduced equation of $p(\bigcup_{H \in \mathcal{A}} H)$, via the identification $\mathcal{O}_V^W \simeq \mathbb{C}[X_1, \dots, X_n]$.

One easily sees that $B(W) \simeq \prod_i B(W_i)$. More generally, all objects studied here behave “semi-simply”, and we may restrict our attention to irreducible complex reflection groups.

Since $B(W)$ is the fundamental group of the complement of an algebraic hypersurface, it is generated by particular elements called *generators-of-the-monodromy* or *meridiens* (see, for example, [18] or [2]). They map to reflections under the natural epimorphism $B(W) \rightarrow W$. The diagrams given in [18] symbolise presentations whose generators are generators-of-the-monodromy (except for the six exceptional types for which no presentation was known).

Definition 1.6. The generators-of-the-monodromy of $B(W)$ are called *braid reflections*.

This terminology was suggested by Broué. It is actually tempting to simply call them *reflections*: since they generate $B(W)$, the braid group appears to be some sort of (infinite) “reflection group”. This guiding intuition is quite effective.

Another natural feature of $B(W)$ is the existence of a natural length function, which is the unique group morphism

$$l : B(W) \rightarrow \mathbb{Z}$$

such that, for all braid reflection $s \in B(W)$, $l(s) = 1$.

Consider the intersection lattice $\mathcal{L}(\mathcal{A}) := \{\bigcap_{H \in \mathcal{A}} H \mid \mathcal{A} \subseteq \mathcal{A}\}$. Elements of $\mathcal{L}(\mathcal{A})$ are called *flats*. It is standard to endow $\mathcal{L}(\mathcal{A})$ with the reversed-inclusion partial ordering:

$$\forall L, L' \in \mathcal{L}(\mathcal{A}), L \leq L' :\Leftrightarrow L \supseteq L'.$$

For $L \in \mathcal{L}(\mathcal{A})$, we denote by L^0 the complement in L of the flats strictly included in L . The $(L^0)_{L \in \mathcal{L}(\mathcal{A})}$ form a stratification $\mathcal{S}$ of V . We consider the partial ordering on $\mathcal{S}$ defined by $L^0 \leq L'^0 :\Leftrightarrow L \leq L'$. This is a degeneracy relation:

$$\forall L \in \mathcal{L}(\mathcal{A}), \overline{L^0} = L = \bigcup_{S \in \mathcal{S}, L^0 \leq S} S.$$

Since W acts on $\mathcal{A}$, it acts on $\mathcal{L}(\mathcal{A})$ and we obtain a quotient stratification $\overline{\mathcal{S}}$ of $W \backslash V$ called *discriminant stratification*.

Proposition 1.7 ([36], Corollary 6.114). *Let $v \in V$. The vectors $\xi_1(v), \dots, \xi_n(v)$ span the tangent space to the stratum of $\mathcal{S}$ containing v . The vectors $\bar{\xi}_1(v), \dots, \bar{\xi}_n(v)$ span the tangent space to the stratum of $\bar{\mathcal{S}}$ containing $\bar{v}$.*

Another chapter of the classical invariant theory of complex reflection groups is Springer's theory of regular elements:

Definition 1.8. Let $W \subseteq \mathrm{GL}(V)$ be a complex reflection group. Let ζ be a complex root of unity. An element $w \in W$ is ζ -regular if $\ker(w - \zeta) \cap V^{\mathrm{reg}} \neq \emptyset$. The eigenvalue ζ is then said to a *regular eigenvalue* and its order d is a *regular number*.

Note that, since W acts freely on V^{reg} , a ζ -regular element must have the same order as ζ . The regularity of ζ only depends on its order d , since the k -th power of a ζ -regular element is ζ^k -regular.

The following theorem compiles some of the main features:

Theorem 1.9. *Let $W \subseteq \mathrm{GL}(V)$ an irreducible complex reflection group, with degrees $d_1, \dots, d_n$ and codegrees $d_1^*, \dots, d_n^*$.*

(1) *Let d be a positive integer. Set*

$$A(d) := \{i = 1, \dots, n \mid d \mid d_i\} \quad \text{and} \quad B(d) := \{i = 1, \dots, n \mid d \mid d_i^*\}.$$

Then $|A(d)| \leq |B(d)|$, and d is regular if and only if $|A(d)| = |B(d)|$.

(2) *Let w be a ζ -regular element of order d . Let $V' := \ker(w - \zeta)$. The centraliser $W' := C_W(w)$, viewed in its natural representation in $\mathrm{GL}(V')$, is a complex reflection group with degrees $(d_i)_{i \in A(d)}$ and codegrees $(d_i^*)_{i \in B(d)}$.*

(3) *Let w be ζ -regular element of order d . Then $W' \backslash V' \simeq (W \backslash V)^{\mu_d}$ and $W' \backslash V'^{\mathrm{reg}} \simeq (W \backslash V^{\mathrm{reg}})^{\mu_d}$ (where the μ_d -action is the quotient action of the scalar multiplication on V).*

Statement (2) was proved by Springer in his seminal paper [41]. Statement (3) was proved independently by Lehrer and Denef-Loeser. Statement (1) was initially observed by Lehrer-Springer on a case-by-case basis, a conceptual proof was recently given by Lehrer-Michel, [31].

Example 1.10. Let $W := G_{37} = W(E_8)$. It is a well-generated complex reflection group in $\mathrm{GL}_8(\mathbb{C})$, whose degrees are

$$2, 8, 12, 14, 18, 20, 24, 30.$$

By duality, the codegrees are

$$0, 6, 10, 12, 16, 18, 22, 28.$$

The integer 4 is regular. The centraliser W' is a complex reflection group of type G_{31} . Its degrees are

$$8, 12, 20, 24,$$

while the codegrees are

$$0, 12, 16, 28.$$

It is not a duality group, and it is not well-generated (see next section).

2. WELL-GENERATED COMPLEX REFLECTION GROUPS

Irreducible complex reflection groups were classified fifty years ago by Shephard and Todd, [42]. There is an infinite family $G(de, e, n)$, where d, e, n are positive integers, and 34 exceptions $G_4, \dots, G_{37}$. Let us distinguish three subclasses of complex reflection groups:

- (complexified) real reflection groups, obtained by scalar extension from reflection groups of real vector spaces;
- 2-reflection groups, generated by reflections of order 2;

- *well-generated reflection groups*, complex reflection groups $W \subseteq \mathrm{GL}(V)$ which can be generated by $\dim_{\mathbb{C}} V/V^W$ reflections, where $V^W := \{v \in V \mid \forall w \in W, wv = v\}$.

Real reflection groups are both 2-reflection groups and well-generated. For non-real groups, any combination of the other two properties may hold.

As far as the $K(\pi, 1)$ conjecture and properties of braid groups are concerned, it is enough to restrict one's attention to 2-reflection groups:

Definition 2.1. Let $W \subseteq \mathrm{GL}(V)$ and $W' \subseteq \mathrm{GL}(V')$ be complex reflection groups. We say that W and W' are *isodiscriminantal* if one may find systems of basic invariants f (resp. f') for W (resp. W') such that $\Delta(W, f) = \Delta(W', f')$.

When this happens, $W \setminus V^{\mathrm{reg}} \simeq W' \setminus V'^{\mathrm{reg}}$ and $B(W) \simeq B(W')$.

Theorem 2.2. *Any complex reflection group is isodiscriminantal to a complex 2-reflection group.*

Proof. This may be observed on the classification and was certainly known to experts. $\square$

Remark 2.3. The recent work of Couwenberg-Heckman-Looijenga, [22], can probably be adapted to provide a direct argument. Let us sketch how one may try to proceed. All references and notations are from [22]. Assume that W is not a 2-reflection group. For each $H \in \mathcal{A}$, let e_H be the order of the pointwise stabiliser W_H and set $\kappa_H := 1 - e_H/2$. Consider the Dunkl connection ∇ with connection form $\sum_{H \in \mathcal{A}} \omega_H \otimes \kappa_H \pi_H$, as in Example 2.5. Since $e_H \geq 2$, we have $\kappa_H \leq 0$. In particular, $\kappa_0 = 1/n \sum_{H \in \mathcal{A}} \kappa_H \leq 0$ (Lemma 2.13) and we are in the situation of *loc. cit.* Section 5. In many cases, this suffices to conclude. The problem is that, even though at least some e_H have to be > 2 , it is possible that $\mathcal{A}$ contains several orbits, some of them with $e_H = 2$. To handle this, one has to enlarge the “Schwarz symmetry group” of *loc. cit.*, Section 4.

The importance of the distinction between well-generated and “badly-generated” groups was first pointed out by Orlik-Solomon, who observed in [34] a coincidence with invariant-theoretical aspects. Their observations may be refined and completed as follows:

Theorem 2.4. *Let W be an irreducible complex reflection group. The following assertions are equivalent:*

- (i) W is well-generated.
- (ii) For all $i \in \{1, \dots, n\}$, $d_i + d_i^* = d_n$.
- (iii) For all $i \in \{1, \dots, n\}$, $d_i + d_i^* \leq d_n$.
- (iv) For any system of basic invariants f , there exists a system of basic derivations ξ such that the discriminant matrix decomposes as $M = M_0 + X_n M_1$, where M_0, M_1 are matrices with coefficients in $\mathbb{C}[X_1, \dots, X_{n-1}]$ and M_1 is lower triangular with non-zero scalars on the diagonal.
- (v) For any system of basic invariants f , we have $\frac{\partial^n \Delta(W, f)}{(\partial X_n)^n} \in \mathbb{C}^\times$ (in other words, $\Delta(W, f)$, viewed as a polynomial in X_n with coefficients in $\mathbb{C}[X_1, \dots, X_{n-1}]$, is monic of degree n).

The matrix M_1 from assertion (iv) is an analogue of the matrix J^* from [39], p. 10. Assertion (iv) itself generalises the non-degeneracy argument for J^* , which is an important piece of the construction of Saito’s “flat structure”.

Proof. (i) $\Rightarrow$ (ii) was observed in [34] inspecting the classification. We still have no good explanation.

(ii) $\Rightarrow$ (iii) is trivial.

(iii) $\Rightarrow$ (v). Let $h := d_n$. A first step is to observe that, under assumption (iii), h is a regular number. Indeed, condition (iii) implies, for any $i = 1, \dots, n-1$, that $0 < d_i < h$ and $0 < d_i^* < h$,

thus that exactly one degree (d_n) and one codegree (d_n^*) are multiples of h , thus h is regular (Theorem 1.9 (1)). Since h is regular and divides only one degree, we may use [2], Lemma 1.6 (ii) to obtain assertion (v): the discriminant is X_n -monic, and by weighted-homogeneity it must be of degree n .

(iii) $\Rightarrow$ (iv) is a refinement of the previous discussion. Each entry $m_{i,j}$ of the matrix M is weighted-homogeneous of weight $d_i + d_j^* \leq d_n + d_1^* = 2h - d_1 < 2h$; since X_n has weight h , $\deg_{X_n} m_{i,j} \leq 1$. This explains the decomposition $M = M_0 + X_n M_1$, where M_0 and M_1 have coefficients in $\mathcal{O}_Y$. If $i < j$ and $d_i < d_j$, then $d_i + d_j^* < d_j + d_j^* = h$, thus $\deg_{X_n} m_{i,j} = 0$. The matrix M_0 is *almost lower triangular*, i.e., lower triangular except that there could be non-zero terms above the diagonal in square diagonal blocks corresponding to successive equal degrees (successive degrees may indeed be equal, as in the example of type D_4 , where the degrees are 2, 4, 4, 6).

Let $i_0 < j_0$ such that $d_{i_0} = d_{i_0+1} = \dots = d_{j_0}$ (looking at the classification, one may observe that this forces $j = i + 1$; this observation is not used in the argument below). For all $i, j \in \{i_0, \dots, j_0\}$, we have $d_i + d_j^* = h$. By weighted homogeneity, this implies that the corresponding square block of M_1 consists of scalars. The basic derivations $\xi_{i_0}, \dots, \xi_{j_0}$ all have the same degrees, thus one is allowed to perform Gaussian elimination on the corresponding columns of M . Thus, up to replacing ξ by another system of basic derivations ξ' , we may assume that M_1 is lower triangular.

The diagonal terms of M_1 must be scalars, once again by weighted homogeneity. Assuming (iii), we already know that (v) holds. The determinant of M is $\Delta(X, f)$; (v) implies that the coefficient of X_n^n is non-zero. This coefficient is the product of the diagonal terms of M_1 . We have proved (iv).

(iv) $\Rightarrow$ (v) is trivial.

(v) $\Rightarrow$ (i) follows from the main result in [2]. □

The following notion was considered in [38] for real reflection groups.

Definition 2.5. A system of basic derivations is *flat* (with respect to f) if the discriminant matrix may be written $M = M_0 + X_n \text{Id}$, where M_0 is a matrix with coefficients in $\mathbb{C}[X_1, \dots, X_{n-1}]$.

Corollary 2.6. *Let W be a well-generated irreducible reflection group, together with a system of basic invariants f . There exists a flat system of basic derivations.*

Proof. Let ξ be any system of basic derivations. Write $M = M_0 + X_n M_1$, as in characterisation (iv) from the above theorem. The matrix M_1 is invertible in $\text{GL}_n(\mathbb{C}[X_1, \dots, X_{n-1}])$. The matrix $M_1^{-1}M = M_1^{-1}M_0 + X_n \text{Id}$ represents a flat system of basic derivations (weighted homogeneity is preserved by the Gaussian elimination procedure). □

Contrary to what happens with real reflection groups, we may not use the identification $V \simeq V^*$ to obtain a “flat system of basic invariants”.

Irreducible groups which are not well-generated may always be generated by $\dim V + 1$ reflections. This fact has been observed long ago, by case-by-case inspection, but no general argument is known. In some sense, these badly-generated groups should be thought of as affine groups. The simplest example of a non-well-generated group is the group $G(4, 2, 2)$, generated by

$$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}.$$

Among high-dimensional exceptional complex reflection groups, only G_{31} is badly-generated (see Example 1.10).

2-reflection groups		Other well-generated groups
Real series	$G(1, 1, n)$ (A_{n-1})	$G_4, G_8, G_{16}, G_{25}, G_{32}$
	$G(2, 1, n)$ (B_{n-1})	$G(d, 1, n), G_5, G_{10}, G_{18}, G_{26}$
Real/complex series	$G(e, e, n)$ (contains D_n and $I_2(e)$)	$G_6, G_9, G_{14}, G_{17}, G_{20}, G_{21}$
Real exceptions	G_{23} (H_3), G_{28} (F_4), G_{30} (F_4) G_{35} (E_6), G_{36} (E_7), G_{37} (E_8)	
Complex exceptions	$G_{24}, G_{27}, G_{29}, G_{33}, G_{34}$	

TABLE 1. Irreducible well-generated complex reflection groups.

In the sequel, several arguments are case-by-case. The list of irreducible well-generated complex reflection groups is given in Table 1. The actual number of cases to consider depends on the type of result:

- Thanks to Theorem 2.2, for any statement about $W \setminus V^{\text{reg}}$ and its topology, one may restrict one's attention to groups generated by involutive reflections, which are listed in the first column of the table.
- Some results involve the actual structure of W (e.g., Proposition 7.5), and groups with higher order reflections have to be considered. These cases are listed in the second column of the table, on the same line as the corresponding 2-reflection groups (see Theorem 2.2).

Lemma 2.7. *Let $W \subseteq \text{GL}(V)$ be a well-generated complex reflection group. Let $v \in V$. Let*

$$V_v := \bigcap_{H \in \mathcal{A}, v \in H} H,$$

$$W_v := \{w \in W \mid wv = v\}.$$

Then W_v may be generated by $\dim_{\mathbb{C}} V/V_v$ reflections. In particular, W_v is again a well-generated complex reflection group.

Proof. The fact that W_v is again a complex reflection group is a classical theorem due to Steinberg. The fact that W_v is again well-generated is easy to check on the classification (it follows for example from Broué-Malle-Rouquier's observation that their diagrams in [18] provide generating systems for representatives of all conjugacy classes of W_v ; when W is real, no case-by-case is needed, since W_v is again real thus well-generated). $\square$

3. SYMMETRIC GROUPS, CONFIGURATIONS SPACES AND CLASSICAL BRAID GROUPS

This section introduces some basic terminology and notations. Everything here is classical and elementary.

Let n be a positive integer. The symmetric group $\mathfrak{S}_n$ may be viewed as reflection group, acting on $\mathbb{C}^n$ by permuting the canonical basis. This representation is not irreducible. Let H be the hyperplane of equation $\sum_{i=1}^n X_i$ (where $X_1, \dots, X_n$ is the dual canonical basis of $\mathbb{C}^n$). It is preserved by $\mathfrak{S}_n$, which acts on it as an irreducible complex reflection group.

We have $\mathbb{C}[X_1, \dots, X_n]^{\mathfrak{S}_n} = \mathbb{C}[\sigma_1, \dots, \sigma_n]$ and $\mathcal{O}_H^{\mathfrak{S}_n} = \mathbb{C}[\sigma_1, \dots, \sigma_n]/\sigma_1$, where $\sigma_1, \dots, \sigma_n$ are the elementary symmetric functions on $X_1, \dots, X_n$. Set $E'_n := \mathfrak{S}_n \setminus \mathbb{C}^n = \text{Spec } \mathbb{C}[\sigma_1, \dots, \sigma_n]$ and $E_n := \mathfrak{S}_n \setminus H = \text{Spec } \mathbb{C}[\sigma_1, \dots, \sigma_n]/\sigma_1$. These spaces have more convenient descriptions in terms of multisets.

Recall that a *multiset* is a set S (the *support* of the multiset) together with a map $m : S \rightarrow \mathbb{Z}_{\geq 1}$ (the *multiplicity*). The *cardinal* of such a multiset is $\sum_{s \in S} m(s)$ (it lies in $\mathbb{Z}_{\geq 0} \cup \{\infty\}$). If (S, m)

and (S', m') are two multisets and if S, S' are subsets of a common ambient set, then we may define a *multiset (disjoint) union* $(S, m) \cup (S', m')$, whose support is $S \cup S'$ and whose multiplicity is $m + m'$ (where m , resp. m' , is extended by 0 outside S , resp. S'). If $(s_1, \dots, s_n)$ is a sequence of elements of a given set S , we use the notation $\{s_1, \dots, s_n\}$ (with brackets in bold font) to refer to the multiset $\bigcup_{i=1}^n (\{s_i\}, 1)$, *i.e.*, the multiset consisting of the s_i 's “taken with multiplicities”.

Let $(x_1, \dots, x_n) \in \mathbb{C}^n$. The associated $\mathfrak{S}_n$ -orbit is uniquely determined by $\{x_1, \dots, x_n\}$. This identifies E'_n with the set of multisets of cardinal n with support in $\mathbb{C}$ (such multisets are called *configurations of n points in $\mathbb{C}$*). The subvariety E_n , defined by $\sigma_1 = 0$, consists of *centered configurations*, *i.e.*, configurations $\{x_1, \dots, x_n\}$ satisfying $\sum_{i=1}^n x_i = 0$. The natural inclusion $E_n \subseteq E'_n$ admits the retraction ρ defined by

$$\rho(\{x_1, \dots, x_n\}) := \{x_1 - \sum_{i=1}^n x_i/n, \dots, x_n - \sum_{i=1}^n x_i/n\}.$$

Algebraically, this corresponds to the identification of $\mathbb{C}[\sigma_1, \dots, \sigma_n]/\sigma_1$ with $\mathbb{C}[\sigma_2, \dots, \sigma_n]$.

We find it convenient to use configurations in E'_n to represent elements of E_n , implicitly working through ρ . *E.g.*, in the proof of Proposition 9.3, it makes sense to describe a deformation retraction of a subspace of E_n to a point in terms of arbitrary configurations because the construction, which only implies the relative values of the x_i 's, is compatible with ρ . We adopt this viewpoint from now on, without further justifications (compatibility will always be obvious).

Consider the lexicographic total ordering of $\mathbb{C}$: if $z, z' \in \mathbb{C}$, we set

$$z \leq z' :\Leftrightarrow \begin{cases} \operatorname{re}(z) < \operatorname{re}(z') & \text{or} \\ \operatorname{re}(z) = \operatorname{re}(z') \text{ and } \operatorname{im}(z) \leq \operatorname{im}(z'). \end{cases}$$

Definition 3.1. The *ordered support* of an element of E_n is the unique sequence $(x_1, \dots, x_k)$ such that the set $\{x_1, \dots, x_k\}$ is the support and $x_1 < x_2 < \dots < x_k$.

We may uniquely represent an element of E_n by its ordered support $(x_1, \dots, x_k)$ and the sequence $(n_1, \dots, n_k)$ of multiplicities at $x_1, \dots, x_k$.

The regular orbit space $E_n^{\text{reg}} := \mathfrak{S}_n \backslash H^{\text{reg}}$ consists of those multisets whose support has cardinal n (or, equivalently, whose multiplicity is constantly equal to 1). More generally, the strata of the discriminant stratification of E_n are indexed by partitions of n : the stratum S_λ associated with a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$, where the λ_j 's are integers with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k > 0$ and $\sum_{j=1}^k \lambda_j = n$, consists of configurations whose supports have cardinal k and whose multiplicity functions take the values $\lambda_1, \dots, \lambda_k$ (with multiplicities).

The braid group B_n associated with $\mathfrak{S}_n$ is the usual braid group on n strings. We need to be more precise about our choice of basepoint. For this purpose, we define

$$E_n^{\text{gen}}$$

as the subset of E_n^{reg} consisting of configurations of n points with distinct real parts (this is the first in a series of definitions of semi-algebraic nature). It is clear that:


Lemma 3.2. E_n^{gen} is contractible.

Using our topological conventions, we set

$$B_n := \pi_1(E_n^{\text{reg}}, E_n^{\text{gen}}).$$

This group admits a *standard generating set* (the one considered by Artin), consisting of braid reflections $\sigma_1, \dots, \sigma_{n-1}$ defined as follows (we used bold fonts to avoid confusion with the elementary symmetric functions). Let $(x_1, \dots, x_n)$ be the ordered support of a point in E_n^{gen} . Then

σ_i is represented by the following motion of the support:



Artin's presentation for B_n is

$$B_n = \langle \sigma_1, \dots, \sigma_{n-1} \mid \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \sigma_i \sigma_j = \sigma_j \sigma_i \text{ if } |i - j| > 1 \rangle.$$

The following definition requires a compatibility condition which is a classical elementary consequence of the above presentation.

Definition 3.3. Let G be a group. The (right) Hurwitz action of B_n on G^n is defined by

$$(g_1, \dots, g_{i-1}, g_i, g_{i+1}, g_{i+2}, \dots, g_n) \cdot \sigma_i := (g_1, \dots, g_{i-1}, g_{i+1}, g_{i+1}^{-1} g_i g_{i+1}, g_{i+2}, \dots, g_n),$$

for all $(g_1, \dots, g_n) \in G^n$ and all $i \in \{1, \dots, n-1\}$.

This action preserves the fibers of the product map $G^n \rightarrow G, (g_1, \dots, g_n) \mapsto g_1 \dots g_n$.

4. AFFINE VAN KAMPEN METHOD

This section contains some generalities about Zariski-Van Kampen techniques. Let $P \in \mathbb{C}[X_1, \dots, X_n]$ be a reduced polynomial. What we have in mind is that P is the discriminant of a complex reflection group, but it does not cost more to work in a general context. Let $\mathcal{H}$ be the hypersurface of $\mathbb{C}^n$ defined by $P = 0$. Van Kampen's method is a strategy for computing a presentation of

$$\pi_1(\mathbb{C}^n - \mathcal{H}).$$

We assume that P actually involves X_n and write

$$P = \alpha_0 X_n^d + \alpha_1 X_n^{d-1} + \alpha_2 X_n^{d-2} + \dots + \alpha_d,$$

where d is a positive integer (the *degree in X_n*) and $\alpha_0, \dots, \alpha_d \in \mathbb{C}[X_1, \dots, X_{n-1}]$, with $\alpha_0 \neq 0$. We say that P is X_n -*monic* if α_0 is a scalar; when one is only interested in the hypersurface defined by P , it is convenient to then renormalise P to have $\alpha_0 = 1$. We denote by $\text{Disc}_{X_n}(P)$ the *discriminant* of P with respect to X_n , i.e., the resultant of P and $\frac{\partial P}{\partial X_n}$.

This discriminant is a non-zero element of $\mathbb{C}[X_1, \dots, X_{n-1}]$ and defines an hypersurface $\mathcal{K}$ in $Y := \text{Spec } \mathbb{C}[X_1, \dots, X_{n-1}] \simeq \mathbb{C}^{n-1}$. Let p be the natural projection $\text{Spec } \mathbb{C}[X_1, \dots, X_n] \twoheadrightarrow Y$.

Definition 4.1. The *bifurcation locus* of P (with respect to the projection p) is the algebraic hypersurface $\mathcal{K} \subseteq Y$ defined by the equation $\text{Disc}_{X_n}(P) = 0$.

Definition 4.2. A point $y \in Y$ is said to be *generic* if it is not in $\mathcal{K}$; a *generic line of direction X_n* is the fiber L_y of p over a generic point $y \in Y$.

We represent points of $\mathbb{C}^n$ by pairs (x, y) where x is the value of the X_n coordinate and y is the image of the point under p . Let $E := p^{-1}(Y - \mathcal{K}) \cap (\mathbb{C}^n - \mathcal{H})$. The projection p restricts to a locally trivial fibration $E \rightarrow Y - \mathcal{K}$, whose fibers are complex lines with d points removed. Choose a basepoint $(x, y) \in E$, let F be the fiber containing (x, y) .

The following basic lemma was brought to my attention by Deligne and should certainly have been included in my earlier paper [2].

Lemma 4.3. *If P is X_n -monic, the fibration $p : E \rightarrow Y - \mathcal{K}$ is split.*

Proof. Assume $\alpha_0 = 1$. Consider $\phi : Y \rightarrow \mathbb{C}, y \mapsto 1 + \sum_{i=1}^d |\alpha_i(y)|$. Then the map $Y - \mathcal{K} \rightarrow E, y \mapsto (\phi(y), y)$ is a splitting. $\square$

As a consequence, the fibration long exact sequence breaks into split short exact sequences. Consider the commutative diagram:

$$\begin{array}{ccccccc}
 1 & \longrightarrow & \pi_1(F, (x, y)) & \xrightarrow{\iota_*} & \pi_1(E, (x, y)) & \xrightarrow{p_*} & \pi_1(Y - \mathcal{K}, y) \longrightarrow 1 \\
 & & & & \downarrow \beta & & \\
 & & & \searrow \alpha & & & \\
 & & & & \pi_1(\mathbb{C}^n - \mathcal{H}, (x, y)). & &
 \end{array}$$

whose first line is a split exact sequence (the end of the fibration exact sequence) and β comes from the inclusion of spaces.

We are in the context of [2, Theorem 2.5], from which we conclude that α is surjective. We can actually be more precise and write a presentation for $\pi_1(\mathbb{C}^n - \mathcal{H}, (x, y))$. The semi-direct structure of $\pi_1(E, (x, y))$ defines a morphism $\Phi : \pi_1(Y - \mathcal{K}, y) \rightarrow \text{Aut}(\pi_1(F, (x, y)))$.

Theorem 4.4 (Van Kampen presentation). *Let $f_1, \dots, f_d$ be generators of $\pi_1(F, (x, y)) \simeq F_d$. Let $g_1, \dots, g_m$ be generators of $\pi_1(Y - \mathcal{K}, y)$ with associated automorphisms $\phi_j := \Phi(g_j)$. We have*

$$\pi_1(\mathbb{C}^n - \mathcal{H}, (x, y)) \simeq \langle f_1, \dots, f_d \mid f_i = \phi_j(f_i), 1 \leq i \leq d, 1 \leq j \leq m \rangle.$$

Proof. Using the semi-direct product structure, we have the presentation

$$\pi_1(E, (x, y)) \simeq \langle f_1, \dots, f_d, g_1, \dots, g_m \mid g_j f_i = \phi_j(f_i) g_j, 1 \leq i \leq d, 1 \leq j \leq m \rangle.$$

One concludes by observing that $g_1, \dots, g_m$ may be chosen to be meridiens (“generators-of-the-monodromy”) around the irreducible components of $p^{-1}(\mathcal{K})$: by [2, Lemma 2.1.(ii)], $\ker \beta$ is generated as a normal subgroup by those meridiens. $\square$

Corollary 4.5 (Explicit Zariski 2-plane section). *Let P be a reduced polynomial in $\mathbb{C}[X_1, \dots, X_n]$. Let $a_1, \dots, a_{n-2} \in \mathbb{C}$. Assume that:*

- (i) P is X_n -monic,
- (ii) the coefficients of $\text{Disc}_{X_n}(P)$ viewed as polynomial with variable X_{n-1} and coefficients in $\mathbb{C}[X_1, \dots, X_{n-2}]$ are altogether coprime (this in particular holds when $\text{Disc}_{X_n}(P)$ is X_{n-1} -monic),
- (iii) $\text{Disc}_{X_{n-1}}(\text{Disc}_{X_n}(P))(a_1, \dots, a_{n-2}) \neq 0$.

Let $\mathcal{H}$ be the hypersurface of $\mathbb{C}^n$ with equation $P = 0$, let Π be the affine 2-plane in $\mathbb{C}^n$ defined by $X_1 = a_1, \dots, X_{n-2} = a_{n-2}$. Then the map $\Pi \cap (\mathbb{C}^n - \mathcal{H}) \hookrightarrow \mathbb{C}^n - \mathcal{H}$ is a π_1 -isomorphism.

Proof. Consider the affine line $p(\Pi) \subseteq Y$. Condition (iii) expresses that $p(\Pi)$ is a generic line of direction X_{n-1} in Y and, under condition (ii), [2, Theorem 2.5], the injection $p(\Pi) \cap (Y - \mathcal{K}) \hookrightarrow Y - \mathcal{K}$ is π_1 -surjective. In particular, in Theorem 4.4, we can choose loops drawn in $\Pi \cap E$ to represent the lifted generators of $\pi_1(Y - \mathcal{K})$. $\square$

Combined with our earlier work with Jean Michel, this corollary suffices to prove Theorem 0.6, since the 2-plane sections described in [8] satisfy conditions (i)–(iii). Presentations for $\pi_1(\Pi \cap (\mathbb{C}^n - \mathcal{H}))$ were obtained in [8], using our software package VKCURVE. At this stage, Theorem 0.6 relies on brutal computations. The sequel will provide a much more satisfying approach.

5. LYASHKO-LOOIJENGA COVERINGS

Let W be an irreducible well-generated complex reflection group, together with a system of basic invariants f and a flat system of basic derivations ξ with discriminant matrix $M_0 + X_n \text{Id}$. Expanding the determinant, we observe that

$$\Delta_f = \det(M_0 + X_n \text{Id}) = X_n^n + \alpha_2 X_n^{n-2} + \alpha_3 X_n^{n-3} + \dots + \alpha_n,$$

where $\alpha_i \in \mathbb{C}[X_1, \dots, X_{n-1}]$. Since Δ_f is weighted homogeneous of total weight nh for the system of weights $\text{wt}(X_i) = d_i$, each α_i is weighted homogeneous of weight ih .

Definition 5.1. The (generalised) *Lyashko-Looijenga morphism* is the morphism LL from $Y = \text{Spec } \mathbb{C}[X_1, \dots, X_{n-1}]$ to $E_n \simeq \text{Spec } \mathbb{C}[\sigma_2, \dots, \sigma_n]$ defined by $\sigma_i \mapsto (-1)^i \alpha_i$.

This is of course much better understood in the following geometric terms. Let $v \in V$. The orbit $\bar{v} \in W \backslash V$ is represented by a pair

$$(x, y) \in \mathbb{C} \times Y,$$

where $x = f_n(v)$ and y is the point in Y with coordinates $(f_1(v), \dots, f_{n-1}(v))$. This coding of points in $W \backslash V$ will be used throughout this article. As in the previous section, we study the space $W \backslash V \simeq \mathbb{C} \times Y$ according to the fibers of the the projection $p : W \backslash V \rightarrow Y, (x, y) \mapsto y$.

Definition 5.2. For any point y in Y , we denote by L_y the fiber of the projection $p : W \backslash V \rightarrow Y$ over y .

For any $y \in Y$, the affine line L_y intersects the discriminant $\mathcal{H}$ in n points (counted with multiplicities), whose coordinates are

$$(x_1, y), \dots, (x_n, y),$$

where $\{x_1, \dots, x_n\}$ is the multiset of solutions in X_n of $\Delta_f = 0$ where each α_i has been replaced by its value at y . We have

$$LL(y) = \{x_1, \dots, x_n\}.$$

The bifurcation locus $\mathcal{K} \subseteq Y$ (see previous section) corresponds precisely to those y such that $LL(y)$ contains multiple points.

The main theorem of this section generalises earlier results from [32]:

Theorem 5.3. *The polynomials $\alpha_2, \dots, \alpha_n \in \mathbb{C}[X_1, \dots, X_{n-1}]$ are algebraically independent and $\mathbb{C}[X_1, \dots, X_{n-1}]$ is a free graded $\mathbb{C}[\alpha_2, \dots, \alpha_n]$ -module of rank $n!h^n/|W|$. As a consequence, LL is a finite morphism. It restricts to an unramified covering $Y - \mathcal{K} \rightarrow E_n^{\text{reg}}$ of degree $n!h^n/|W|$.*

I thank Eduard Looijenga for precious help with the theorem. A prior version of this text contained a gap (the key Lemma 5.5) and the very nice argument below is due to him.

Lemma 5.4. *Let $v \in V$ with image $\bar{v} \in W \backslash V$. Let*

$$V_v := \bigcap_{H \in \mathcal{A}, v \in H} H.$$

The multiplicity of $\mathcal{H}$ at $\bar{v}$ is $\dim_{\mathbb{C}} V/V_v$.

Proof. At $v = 0$, the multiplicity is the valuation of Δ_f , which is indeed n .

When $v \neq 0$, we consider the parabolic subgroup $W_v := \{w \in W | wv = v\}$. By Lemma 2.7, W_v is again a well-generated complex reflection group. The quotient map $\bigcup_{H \in \mathcal{A}} H \rightarrow \mathcal{H} = W \backslash (\bigcup_{H \in \mathcal{A}} H)$ factors through $\bigcup_{H \in \mathcal{A}} H \rightarrow W_v \backslash (\bigcup_{H \in \mathcal{A}} H)$. Because $W_v \backslash (\bigcup_{H \in \mathcal{A}} H) \rightarrow W \backslash (\bigcup_{H \in \mathcal{A}} H)$ is unramified over $\bar{v}$, the multiplicity of $\mathcal{H}$ at $\bar{v}$ coincides with the multiplicity of $W_v \backslash (\bigcup_{H \in \mathcal{A}} H)$ at the image $\tilde{v}$ of v . Around $\tilde{v}$, $W_v \backslash (\bigcup_{H \in \mathcal{A}} H)$ is the same as $W_v \backslash (\bigcup_{H \in \mathcal{A}, v \in H} H)$. This hypersurface is a direct product of V_v with the discriminant of W_v . The multiplicity at $\tilde{v}$ is the multiplicity at the origin of the discriminant of W_v . After reduction to the irreducible case, we apply the already solved case: the multiplicity is the rank of W_v or, in other words, $\dim_{\mathbb{C}} V/V_v$. $\square$

Lemma 5.5. $LL^{-1}(0) = \{0\}$

Proof. Let C be the tangent cone to $\mathcal{H}$ in $\mathbb{C}^n \simeq \mathbb{C} \times Y$. It is a closed subvariety of the tangent bundle to $\mathbb{C}^n$. Because $\mathcal{H}$ is quasi-homogeneous and $\forall i \in \{1, \dots, n-1\}, \text{wt}(X_n) > \text{wt}(X_i)$, the cone C is “horizontal” at 0. In particular, the (fibre over 0 of the tangent cone to the) “vertical” line L_0 of equation $X_1 = \dots = X_{n-1} = 0$ is not in C .

Let $y = (x_1, \dots, x_{n-1}) \in LL^{-1}(0)$, i.e., such that the line L_y with $X_1 = x_1, \dots, X_{n-1} = x_{n-1}$ intersects $\mathcal{H}$ in only one point, $(0, y) \in \mathbb{C} \times Y$. We want to prove that $y = 0$. Because $\mathcal{H}$ is quasi-homogeneous, it is enough to work in a neighborhood of the origin. In particular, we may assume that y is close enough to 0 for L_y to still be outside C . Using a refined Bezout theorem ([29, Corollary 12.4]), we have

$$i((0, y), L_y \cdot \mathcal{H}; \mathbb{C}^n) = m_{(0, y)}(\mathcal{H})$$

where

- $i((0, y), L_y \cdot \mathcal{H}; \mathbb{C}^n)$ is the intersection multiplicity of $\mathcal{H}$ and L_y at $(0, y)$. This is the order of 0 as a root of the polynomial $\Delta_f|_{X_1=x_1, \dots, X_{n-1}=x_{n-1}}$. By assumption, this is n .
- $m_{(0, y)}(\mathcal{H})$ is the multiplicity of $\mathcal{H}$ at $(0, y)$. Let v be a preimage of $(0, y)$ in V . By Lemma 5.4, $m_{(0, y)}(\mathcal{H}) = \dim_{\mathbb{C}} V/V_v$.

Thus $\dim_{\mathbb{C}} V/V_v = n$, $V_v = 0$, $W_v = W$ and $v = 0$. $\square$

Because LL is quasi-homogeneous, Lemma 5.5 implies that LL is a finite (= quasi-finite and proper) morphism, or, in other words, that $\mathbb{C}[X_1, \dots, X_{n-1}]$ is a finite graded $\mathbb{C}[\alpha_2, \dots, \alpha_n]$ -module. In particular, $\alpha_2, \dots, \alpha_n$ are algebraically independent.

Because $\mathbb{C}[X_1, \dots, X_{n-1}]$ is Cohen-Macaulay and is finite over $\mathbb{C}[\alpha_2, \dots, \alpha_n]$, it is a free $\mathbb{C}[\alpha_2, \dots, \alpha_n]$ -module. The rank may be computed by comparing Hilbert series. Since each X_i has weight d_i , the Hilbert series of $\mathbb{C}[X_1, \dots, X_{n-1}]$ is $\prod_{i=1}^{n-1} \frac{1}{1-t^{d_i}}$. Since each α_i has weight ih , the Hilbert series of $\mathbb{C}[\alpha_2, \dots, \alpha_n]$ is $\prod_{i=2}^n \frac{1}{1-t^{ih}}$. The rank is

$$\prod_{i=1}^{n-1} \frac{(i+1)h}{d_i} = \frac{n!h^{n-1}}{d_1 \dots d_{n-1}} = \frac{n!h^n}{|W|}.$$

Theorem 5.3 now follows from the following generalisation of [32, Theorem 1.4]:

Lemma 5.6. *LL is étale on $Y - \mathcal{K}$.*

Proof. As mentioned in [32, (1.5)], the result will follow if we prove that for all $y \in Y - \mathcal{K}$ with $LL(y) = \{x_1, \dots, x_n\}$, the hyperplanes $H_1, \dots, H_n$ tangent to $\mathcal{H}$ at the n distinct points $(x_1, y), \dots, (x_n, y)$ are in general position.

To prove this, we use Proposition 1.7. Each H_i is spanned by $\bar{\xi}_1(x_i, y), \dots, \bar{\xi}_n(x_i, y)$. Let $(\varepsilon_1, \dots, \varepsilon_n)$ be the basis of $(W \setminus V)^*$ dual to $(\frac{\partial}{\partial X_1}, \dots, \frac{\partial}{\partial X_n})$. Let $l_i = \sum_j \lambda_{j,i} \varepsilon_j$ be a non-zero vector in $(W \setminus V)^*$ orthogonal to H_i . This amounts to taking a non-zero column vector $(\lambda_{j,i})_{j=1, \dots, n}$ in the kernel of $M(x_i, y)$, or equivalently an eigenvector of $M_0(y)$ associated to the eigenvalue $-x_i$. By assumption, the x_i are distinct. The eigenvectors are linearly independent. $\square$

The theorem has the following corollary (which we won't use).

Corollary 5.7. *The space $Y - \mathcal{K}$ is a $K(\pi, 1)$.*

For the sake of clarity, let us also mention:

Corollary 5.8. *Let $v \in V$. Denote by $(x, y) \in \mathbb{C} \times Y$ (identified with $W \setminus V$) the image of v . The following integers coincide:*

- (i) *The multiplicity of x in $LL(y)$.*

- (ii) *The intersection multiplicity of L_y with $\mathcal{H}$ at (x, y) .*
- (iii) *The multiplicity of $\mathcal{H}$ at (x, y) .*
- (iv) *The rank $\dim_{\mathbb{C}} V/V_v$ of the parabolic subgroup W_v .*

Proof. The integer defined by (i) and (ii) are the same by their very definition. The identity between (iii) and (iv) is Lemma 5.4. The argument used to prove Lemma 5.5 also shows the identity between (ii) and (iii). $\square$


The discriminant stratification of E_n yields a natural stratification of Y : when λ is a partition of n , the stratum Y_λ consists of points y such that the multiplicities of $LL(y)$ are distributed according to λ . Applying the corollary, one sees that the stratification $Y = \bigsqcup_{\lambda \models n} Y_\lambda$ is the “shadow” of the discriminant stratification restricted to $\mathcal{H}$.

6. TUNNELS, LABELS AND HURWITZ RULE

Let W be an irreducible well-generated complex reflection group. We keep the notations from the previous section. Let $y \in Y$. Let U_y be the complement in L_y of the vertical imaginary half-lines below the points of $LL(y)$, or in more formal terms:

$$U_y := \{(z, y) \in L_y \mid \forall x \in LL(y), \text{re}(z) = \text{re}(x) \Rightarrow \text{im}(z) > \text{im}(x)\}.$$

Here is an example where the support of $LL(y)$ consists of 4 points, and U_y is the complement of three half-lines:



“Generically”, U_y is the complement of n vertical half-lines. We have to be careful about what “generically” means here: a prerequisite is that $LL(y)$ should consist of n distinct points, which amounts to $y \in Y - \mathcal{K}$ or equivalently

$$LL(y) \in E_n^{\text{reg}},$$

but this is not enough: one needs these points to be on distinct vertical lines, or equivalently that

$$LL(y) \in E_n^{\text{gen}}.$$

Definition 6.1. We set $Y^{\text{gen}} := LL^{-1}(E_n^{\text{gen}})$.

This space, being the “fiber” of the covering LL over the “basepoint” of the basespace E_n^{reg} , is equipped with a Galois action of $B_n := \pi_1(E_n^{\text{reg}}, E_n^{\text{gen}})$.

Let

$$\mathcal{U} := \bigcup_{y \in Y} U_y.$$

Lemma 6.2. *The subspace $\mathcal{U}$ is dense in $W \setminus V^{\text{reg}}$, open and contractible.*

Proof. The first two statements are clear.

Define a continuous function $\beta : Y \rightarrow \mathbb{R}$ by

$$\beta(y) := \max\{\text{im}(x) \mid x \in LL(y)\} + 1.$$

Points of $W \setminus V$ are represented by pairs $(x, y) \in \mathbb{C} \times Y$, or equivalently by triples $(a, b, y) \in \mathbb{R} \times \mathbb{R} \times Y$, where $a = \text{re}(x)$ and $b = \text{im}(x)$. For $t \in [0, 1]$, define $\phi_t : W \setminus V \rightarrow W \setminus V$ by

$$\phi_t(a, b, y) := \begin{cases} (a, b, y) & \text{if } b \geq \beta(y), \\ (a, b + t(\beta(y) - b), y) & \text{if } b \leq \beta(y). \end{cases}$$

Each ϕ_t preserves $\mathcal{U}$ and the homotopy ϕ restricts to a deformation retraction of $\mathcal{U}$ to

$$\bigcup_{y \in Y} \{(x, y) \in L_y \mid \text{im}(x) \geq \beta(y)\}.$$

The latter is a locally trivial bundle over the contractible space Y , with contractible fibers (the fibers are half-planes). Thus it is contractible. $\square$

As explained in the topological preliminaries, we may (and will) use $\mathcal{U}$ as “basepoint” for $W \setminus V^{\text{reg}}$ and refine our definition of $B(W)$:

Definition 6.3. The *braid group* of W is $B(W) := \pi_1(W \setminus V^{\text{reg}}, \mathcal{U})$.


As for other notions actually depending on W , we often write B instead of $B(W)$, since most of the time we implicitly refer to a given W .

Remark 6.4. We will need to consider a natural projection $\pi : B \rightarrow W$. Recall that such a morphism is part of the fibration exact sequence

$$1 \longrightarrow \pi_1(V^{\text{reg}}) \longrightarrow \pi_1(W \setminus V^{\text{reg}}) \xrightarrow{\pi} W \longrightarrow 1.$$

For this exact sequence to be well-defined, one has to make consistent choices of basepoints in V^{reg} and in $W \setminus V^{\text{reg}}$. We have already described our “basepoint” $\mathcal{U}$ in $W \setminus V^{\text{reg}}$. Choose $u \in \mathcal{U}$ and choose a preimage $\tilde{u}$ of u in V^{reg} . If $u' \in \mathcal{U}$ is another choice and if γ is a path in $\mathcal{U}$ from u to u' , γ lifts to a unique path $\tilde{\gamma}$ starting at $\tilde{u}$; since $\mathcal{U}$ is contractible, the fixed-endpoint homotopy class of $\tilde{\gamma}$ (and, in particular, its final point) does not depend on γ . In other words, once we have chosen a preimage of one point of $\mathcal{U}$, we have a natural section $\tilde{\mathcal{U}}$ of $\mathcal{U}$ in V^{reg} , as well as a transitive system of isomorphisms between $(\pi_1(V^{\text{reg}}, \tilde{u}))_{\tilde{u} \in \tilde{\mathcal{U}}}$. From now on, we assume we have made such a choice, and we define the pure braid group as $\pi_1(V^{\text{reg}}, \tilde{\mathcal{U}})$. This selects one particular morphism π (the $|W|$ possible choices yield conjugate morphisms).

Definition 6.5. A *semitunnel* is a triple $T = (y, z, L) \in Y \times \mathbb{C} \times \mathbb{R}_{\geq 0}$ such that $(z, y) \in \mathcal{U}$ and the affine segment $[(z, y), (z + L, y)]$ lies in $W \setminus V^{\text{reg}}$. The path γ_T associated with T is the path $t \mapsto (z + tL, y)$. The semitunnel T is a *tunnel* if in addition $(z + L, y) \in \mathcal{U}$.



The distinction between tunnels and semitunnels should be understood in light of our topological conventions: if T is a tunnel, γ_T represents an element

$$b_T \in \pi_1(W \setminus V^{\text{reg}}, \mathcal{U}),$$

while semitunnels will be used to represent points of the universal cover $(\widetilde{W \setminus V^{\text{reg}}})_{\mathcal{U}}$ (see Section 10).

Definition 6.6. An element $b \in B$ is *simple* if $b = b_T$ for some tunnel T . The set of simple elements in B is denoted by S .


Each tunnel lives in a given L_y , where it may be represented by an horizontal (= constant imaginary part) segment avoiding $LL(y)$ and with endpoints in U_y . The triple (y, z, L) may be uniquely recovered from $[(z, y), (z + L, y)]$. A frequent abuse of terminology will consist of using the term *tunnel* (or *semitunnel*) to designate either the triple (y, z, L) , or the segment $[(z, y), (z +$

$L, y]$, or the pair $(y, [z, z+L])$, depending on the context (in particular, when intersecting tunnels with geometric objects, the tunnels should be understood as affine segments).


Let $y \in Y$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$. The space $p_x((L_y \cap W \setminus V^{\text{reg}}) - U_y)$ is an union of k disjoint open affine intervals $I_1, \dots, I_k$, where

$$I_i := \begin{cases} (x_i - \sqrt{-1}\infty, x_i) & \text{if } i = 1 \text{ or } (i > 1 \text{ and } \text{re}(x_{i-1}) < \text{re}(x_i)), \\ (x_{i-1}, x_i) & \text{otherwise.} \end{cases}$$

(by $(x_i - \sqrt{-1}\infty, x_i)$, we mean the open vertical half-line below x_i). In the first case (when I_i is not bounded), we say that x_i is *deep*. In the picture below, there are three deep points, x_1 , x_3 and x_4 .



Choose a *system of elementary tunnels* for y . By this, we mean the choice, for each $i = 1, \dots, k$, of a small tunnel T_i in L_y crossing I_i and not crossing the other intervals; let $s_i := b_{T_i}$ be the associated element of B .



These elements depend only on y and not on the explicit choice of elementary tunnels.

Definition 6.7. The sequence $lbl(y) := (s_1, \dots, s_k)$ is the *label* of y . Let $i_1, i_2, \dots, i_l$ be the indices of the successive deep points of $LL(y)$. The *deep label* of y is the subsequence $(s_{i_1}, \dots, s_{i_l})$.

In the above example, the deep label is (s_1, s_3, s_4) . The length of the label is n if and only if $y \in Y - \mathcal{K}$. In this case, the deep label coincides with the label if and only if $y \in Y^{\text{gen}}$.

Later on, it will appear that the pair $(LL(y), lbl(y))$ uniquely determines y (Theorem 7.9).

Remark 6.8. When y is generic, $lbl(y)$ is an n -tuple of braid reflections (because an elementary tunnel crossing the interval below a point in $LL(y)$ is essentially the same as a small circle around this point; when y is generic, the points in $LL(y)$ correspond to smooth points of the discriminant, and the elementary tunnels represent generators of the monodromy).

Consider the case $y = 0$ (given by the equations $X_1 = 0, \dots, X_{n-1} = 0$). The multiset $LL(y)$ has support $\{0\}$ with multiplicity n .

Definition 6.9. We denote by δ the simple element such that $lbl(0) = (\delta)$.

This element plays the role of Deligne's element Δ . Choose $v \in V^{\text{reg}}$ such that the W -orbit $\bar{v}$ lies in L_0 . Broué-Malle-Rouquier consider the element (denoted by π , [18, Notation 2.3]) in the pure braid group $P(W)$ represented by the loop

$$\begin{aligned} [0, 1] &\longrightarrow V^{\text{reg}} \\ t &\longmapsto v \exp(2\sqrt{-1}\pi t). \end{aligned}$$

We prefer a different notation:

Definition 6.10. We call *full-twist* this element of $P(W)$, and denote it by τ .

They observe that this element lies in the centre of B ([18, Theorem 2.24]) and conjecture that it generates the center of P .

Since X_n has weight h , δ^h coincides with τ . More precisely, δ is represented by the loop which is the image in $W \setminus V^{\text{reg}}$ of the path in V^{reg}

$$\begin{aligned} [0, 1] &\longrightarrow V^{\text{reg}} \\ t &\longmapsto v \exp(2\sqrt{-1}\pi t/h) \end{aligned}$$

In particular:

Lemma 6.11. *The element $\tau = \delta^h$ is central in $B(W)$ and lies in $P(W)$. The image of δ in W is ζ_h -regular, in the sense of Springer (see Definition 1.8).*

Remark 6.12. The centre of W is a cyclic group of order $\gcd(d_1, \dots, d_n)$. The above argument can be applied to show that $\delta^{h'} \in ZB(W)$, where $h' := \frac{h}{\gcd(d_1, \dots, d_n)}$. Later on (Section 12), we will show that $ZB(W) = \langle \delta^{h'} \rangle$.

Each tunnel lives in a single fiber L_y . Let us now explain how simple elements represented by tunnels living in different fibers may be compared. The idea is that, since being a tunnel is an open condition, one may perturb y without affecting the simple element:

Definition 6.13. Let $T = (y, z, L)$ be a tunnel. A T -neighbourhood of y is a path-connected neighbourhood Ω of y in Y such that, for all $y' \in \Omega$, $T' := (y', z, L)$ is a tunnel.

Such neighbourhoods clearly exist for all $y \in Y$.

Lemma 6.14 (Hurwitz rule). *Let $T = (y, z, L)$ be a tunnel, representing a simple element s . Let Ω be a T -neighbourhood of y . For all $y' \in \Omega$, $T' := (y', z, L)$ represents s .*

Proof. This simply expresses that the tunnels (y', z, L) and (y, z, L) represent homotopic paths, which is clear by definition of Ω . $\square$

Remark 6.15. Let $T_1, \dots, T_k$ is a system of elementary tunnels for y . Let Ω_i be a T_i -neighbourhood for y . A *standard neighbourhood* of y could be defined as a path-connected neighbourhood Ω of y inside $\cap_{i=1}^k \Omega_i$. These standard neighbourhoods form a basis for the topology of Y . A consequence of Hurwitz rule is that the label of y may uniquely be recovered once we know the label of a single $y' \in \Omega$. The converse is not true, unless $y \in Y - \mathcal{K}$. Let $y' \in \Omega$, let S and S' be the strata of $LL(y)$ and $LL(y')$ (with respect to the discriminant stratification of E_n). We have $S' \leq S$. The label of y' is uniquely determined by the label of y if and only if $S = S'$. Otherwise, the ambiguity is caused by the extra ramification at y .

The remainder of this section consists of various consequences of Hurwitz rule.


Corollary 6.16. *Let $y \in Y$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$, and $(n_1, \dots, n_k)$ be the multiplicities. For any i , the canonical length of s_i is given by*

$$l(s_i) = \sum_{j \text{ s.t. } \text{re}(x_i) = \text{re}(x_j) \text{ and } \text{im}(x_i) \leq \text{im}(x_j)} n_j.$$

Proof. The case $y \in Y^{\text{gen}}$ is a consequence of Remark 6.8. The general case follows by perturbing and applying Hurwitz rule. $\square$

Corollary 6.17. *Let $y \in Y$. Let $(s_{i_1}, \dots, s_{i_l})$ be the deep label of y . We have $s_{i_1} \dots s_{i_l} = \delta$.*

Proof. Any tunnel T deep enough and long enough represents $s_{i_1} \dots s_{i_l}$.




The origin $0 \in Y$ lies in a T -neighbourhood of y . To conclude, apply Hurwitz rule. $\square$

Corollary 6.18. *Let $x \in E_n^{\text{gen}}$. Let $\beta \in \pi_1(E_n^{\text{reg}}, x)$, $y \in LL^{-1}(x)$ and $y' := y \cdot \beta$. Let $(b_1, \dots, b_n)$ be the label of y and $(b'_1, \dots, b'_n)$ be the label of y' . Then*

$$(b'_1, \dots, b'_n) = (b_1, \dots, b_n) \cdot \beta,$$

where β acts by right Hurwitz action.

Proof. It is enough to prove this for a standard generator σ_i . Let $(x_1, \dots, x_n)$ be the ordered support of x . By Hurwitz rule, we may adjust the imaginary parts of the x_i 's without affecting the label; in particular we may assume that $\text{im}(x_i) < \text{im}(x_{i+1})$. We may find tunnels $T_- = (y, z_-, L)$ and $T_+ = (y, z_+, L)$ as in the picture below:



The path in E_n^{reg} where x_{i+1} moves along the dotted arrow and all other points are fixed represents σ_i . Applying Hurwitz rule to T_+ , we obtain $b'_i = b_{i+1}$; applying Hurwitz rule to T_- , we obtain $b'_i b'_{i+1} = b_i b_{i+1}$. The result follows. $\square$

Corollary 6.19. *Let $y \in Y^{\text{gen}}$. The cardinal of the Hurwitz orbit $\text{lbl}(y) \cdot B_n$ is at most $n!h^n/|W|$, and there is an equivalence between:*

- (i) $|\text{lbl}(y) \cdot B_n| = n!h^n/|W|$.
- (ii) The orbits $y \cdot B_n$ and $\text{lbl}(y) \cdot B_n$ are isomorphic as B_n -sets.
- (iii) The map $Y^{\text{gen}} \rightarrow E_n \times B_n, y \mapsto (LL(y), \text{lbl}(y))$ is injective.

In the next section, we will prove that conditions (i)–(iii) actually hold. This is not a trivial statement.

Proof. Let G be a group and Ω, Ω' be two G -sets, together with a G -set morphism $\rho : \Omega \rightarrow \Omega'$. Assume that Ω' is transitive. Then ρ is surjective. Assume in addition that Ω is finite. Then Ω' is finite, $|\Omega'| \leq |\Omega|$ and


$$|\Omega'| = |\Omega| \Leftrightarrow \rho \text{ is injective} \Leftrightarrow \rho \text{ is an isomorphism.}$$

By the previous corollary, one has $\text{lbl}(y \cdot \beta) = \text{lbl}(y) \cdot \beta$. In other words, the map lbl extends to a B_n -set morphism $y \cdot B_n \rightarrow \text{lbl}(y) \cdot B_n$. We apply our discussion to $G := B_n$, $\Omega := y \cdot B_n$ and $\Omega' := \text{lbl}(y) \cdot B_n$. Both B_n -sets are clearly transitive. Since $y \in Y - \mathcal{K}$, we have $|LL^{-1}(LL(y))| = |y \cdot B_n| = n!h^n/|W|$ (Theorem 5.3). We deduce that $|\text{lbl}(y) \cdot B_n| \leq n!h^n/|W|$ and (i) $\Leftrightarrow$ (ii).

Assertion (iii) amounts to saying that, for all $y \in Y^{\text{gen}}$, lbl is injective on the fiber of LL containing y . This fiber is precisely the orbit $y \cdot B_n$. Under this rephrasing, it is clear that (ii) $\Leftrightarrow$ (iii). $\square$

Corollary 6.20. *Let s be a simple element. There exists $y \in Y^{\text{gen}}$ and $i \in \{1, \dots, n\}$ such that $s = s_1 \dots s_i$, where $(s_1, \dots, s_n) := \text{lbl}(y)$.*

Proof. Let T be a tunnel representing s . Any T -neighbourhood of y contains generic points. Up to perturbing y , we may assume that $y \in Y^{\text{gen}}$. The picture below explains, on an example, how to move certain points (following the dotted paths) of the underlying configuration to reach a suitable y' :



This path in E_n^{reg} lifts, via LL , to a path in a T -neighbourhood of y whose final point y' satisfies the conditions of the lemma. $\square$

7. REDUCED DECOMPOSITIONS OF COXETER ELEMENTS

Definition 7.1. Let W be an irreducible well-generated complex reflection group. An element $c \in W$ is a (*generalised*) *Coxeter element* if it is ζ_h -regular. More generally, if W is a well-generated complex reflection group decomposed as a sum $W = \bigoplus_i W_i$ of irreducible groups, a Coxeter element in W is a product $c = \prod_i c_i$ of Coxeter elements in each W_i .

When W is irreducible, we may use the constructions from the previous section and the morphism $\pi : B(W) \twoheadrightarrow W$ to obtain typical Coxeter elements (see Definition 6.9):

Lemma 7.2. *When W is irreducible, the element $c := \pi(\delta)$ is a Coxeter element in W .*

Proof. It is a rephrasing of Lemma 6.11. $\square$

The other Coxeter elements, which are conjugates of c , appear when considering other base-points over $\mathcal{U}$ (see Remark 6.4).

More generally, we have:

Lemma 7.3. *Let $y \in Y$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$, let $(n_1, \dots, n_k)$ be the multiplicities. Assume that $\text{re}(x_1) < \dots < \text{re}(x_k)$. Let $(s_1, \dots, s_k) := \text{lbl}(y)$.*

For all i , set $c_i := \pi(s_i)$. Then there exists a preimage $v_i \in V$ of $(x_i, y) \in \mathbb{C} \times Y \simeq W \setminus V$ such that c_i is a Coxeter element in the parabolic subgroup W_{v_i} .

In particular, if $n_i = 1$, then c_i is a reflection.

Proof. When $n_i = n$ (thus $i = k = 1$), the result is Lemma 7.2.

When $n_i = 1$, as pointed out in Remark 6.8, the element s_i is represented by a small loop around a smooth point in the discriminant, thus is a braid reflection, and maps to a reflection in W .

The general case is similar: locally near (x_i, y) , the discriminant is a direct product of V_{v_i} with the discriminant of W_{v_i} (see the proof of Lemma 5.4). This local structure provides a specific morphism $B(W_{v_i}) \rightarrow B(W)$, such that the element “ δ_i ” in $B(W_{v_i})$ (the product of the δ ’s associated with each irreducible components of W_{v_i}) maps to s_i . The lemma follows. $\square$

The assumption $\text{re}(x_1) < \dots < \text{re}(x_k)$ may be removed at the cost of replacing $\pi(s_i)$ by $\pi(s_{i-1})^{-1}\pi(s_i)$ when $\text{re}(x_{i-1}) = \text{re}(x_i)$.

Let R be the set of all reflections in W . As in [3], for all $w \in W$, we denote by $\text{Red}_R(w)$ the set of *reduced R -decompositions* of w , i.e., minimal length sequences of elements of R with product w . Since R is closed under conjugacy, $\text{Red}_R(c)$ is stable under Hurwitz action. We also consider the length function $l_R : W \rightarrow \mathbb{Z}_{\geq 0}$, whose value at w is the common length of the elements of $\text{Red}_R(w)$, and two partial orderings of W defined as follows: for all $w, w' \in W$, we set

$$w \preceq_R w' :\Leftrightarrow l_R(w) + l_R(w^{-1}w') = l_R(w')$$

and

$$w' \succ_R w :\Leftrightarrow l_R(w'w^{-1}) + l_R(w) = l_R(w').$$

Since R is invariant by conjugacy, we have $w \preccurlyeq_R w' \Leftrightarrow w' \succcurlyeq_R w$.

Let $y \in Y^{\text{gen}}$, let $(s_1, \dots, s_n) := \text{lbl}(y)$, let $(r_1, \dots, r_n) := (\pi(s_1), \dots, \pi(s_n)) = \pi^n(\text{lbl}(y))$. By Lemma 7.3, $(r_1, \dots, r_n) \in \text{Red}_R(c)$.

The key result of this section is:

Theorem 7.4. *Let $y \in Y^{\text{gen}}$, The maps*

$$y \cdot B_n \xrightarrow{\text{lbl}} \text{lbl}(y) \cdot B_n \xrightarrow{\pi^n} \pi^n(\text{lbl}(y)) \cdot B_n$$

are isomorphisms of B_n -sets, where $y \cdot B_n$ is the Galois orbit of y , and $\text{lbl}(y) \cdot B_n$ and $\pi^n(\text{lbl}(y)) \cdot B_n$ are Hurwitz orbits.

The theorem implies that conditions (i)–(iii) from Corollary 6.19 actually hold.

In the real case, this was initially conjectured by Looijenga, [32, (3.5)], and proved in a letter from Deligne to Looijenga (crediting discussions with Tits and Zagier), [26]; an equivalent property ([3, Fact 2.2.4]) was independently used in our earlier construction of the dual braid monoid.

These proofs for the real case are based on case-by-case numerology: because the B_n -sets are transitive, it suffices to prove that the cardinality of $y \cdot B_n$ (which is by construction the degree of LL) coincides with that of $(r_1, \dots, r_n) \cdot B_n$. This is an enumeration problem in W and may be tackled by case-by-case analysis (the infinite family are easy to deal with, computers can take care of the exceptional types).

This enumerative approach carries on to our setting: Theorem 7.4 immediately follows from Proposition 7.5 below.

Proposition 7.5. *Let W be a well-generated complex reflection group. Let c be a Coxeter element in W . The Hurwitz action is transitive on $\text{Red}_R(c)$. When W is irreducible, one has $|\text{Red}_R(c)| = n!h^n/|W|$.*

Proof. The proposition clearly reduces to the case when W is irreducible and $w = c$: in the reducible case, reduced decompositions of Coxeter elements are “shuffles” of reduced decompositions of the Coxeter elements of the irreducible summands.

We prove the result case-by-case (see Table 1 in section 2 for the list of cases to be considered). The complexified real case is studied in [26] (transitivity is easy and does not require case-by-case, see for example [3, Proposition 1.6.1]). The $G(e, e, r)$ case combines two results from [6]: Proposition 6.1 (transitivity) and Theorem 8.1 (cardinality).

The case of $G(d, 1, r)$ goes as follows. For all integers i, j with $1 \leq i < j \leq n$, denote by $\tau_{i,j}$ the permutation matrix associated with the transposition $(i\ j) \in \mathfrak{S}_n = G(1, 1, n) \hookrightarrow G(d, 1, r)$. For all $\zeta \in \mu_d$ and all $i \in \{1, \dots, n\}$, denote by $\rho_{i,\zeta}$ the diagonal matrix $\text{Diag}(1, \dots, 1, \zeta, 1, \dots, 1)$, where ζ is in i -th position. There are two types of reflections in $G(d, 1, n)$: *long* reflections are elements of the form $\rho_{i,\zeta}$, with $\zeta \neq 1$; *short* reflections are elements of the form $\tau_{i,j}^\zeta := \rho_{i,\zeta}^{-1} \tau_{i,j} \rho_{i,\zeta}$, with $\zeta \in \mu_d$ and $1 \leq i < j \leq n$. A typical Coxeter element is $c := \rho_{1,\zeta_d} \tau_{1,2} \tau_{2,3} \dots \tau_{n-1,n}$. Since Coxeter elements form a single conjugacy class, and since R is invariant under conjugacy, it suffices to prove the claims for this particular c . Let $(r_1, \dots, r_n) \in \text{Red}_R(c)$. Let us prove that it is Hurwitz equivalent to $(\rho_{1,\zeta_d}, \tau_{1,2}, \tau_{2,3}, \dots, \tau_{n-1,n})$. Consider the morphism $G(d, 1, n) \rightarrow G(1, 1, n)$, $g \mapsto \bar{g}$ sending a monomial matrix to the underlying permutation matrix. This map sends $\tau_{i,j}^\zeta$ to $\tau_{i,j}$ and $\rho_{i,\zeta}$ to 1. The element $\bar{c}$ is a Coxeter element in $G(1, 1, n)$. One deduces that there is a unique long reflection r_{i_0} among $r_1, \dots, r_n$ and that $(r_1, \dots, \widehat{r_{i_0}}, \dots, r_n)$ is a reduced R -decomposition of $\bar{c}$ in $G(1, 1, n)$. Up to applying suitable Hurwitz moves, we may assume

that $i_0 = 1$. Using the transitivity result already known in the type $G(1, 1, n)$ case, we see that $(r_1, \dots, r_n)$ is Hurwitz equivalent to $(\rho_{i,\zeta}, \tau_{1,2}^{\alpha_1}, \dots, \tau_{n-1,n}^{\alpha_{n-1}})$, where $i \in \{1, \dots, n\}$, $\zeta \in \mu_e - \{1\}$ and $\alpha_1, \dots, \alpha_{n-1} \in \mu_d$. By considering the determinant, we see that $\zeta = \zeta_d$. A direct computation shows that, if $i > 1$,

$$\rho_{i,\zeta} \tau_{1,2}^{\alpha_1} \dots \tau_{i-1,i}^{\alpha_{i-1}} = \tau_{1,2}^{\alpha_1} \dots \tau_{i-1,i}^{\alpha_{i-1}} \rho_{i-1,\zeta}.$$

One may use this relation to construct an explicit sequence of Hurwitz moves showing that $(r_1, \dots, r_n)$ is equivalent to $(\rho_{1,\zeta_d}, \tau_{1,2}^{\alpha'_1}, \dots, \tau_{n-1,n}^{\alpha'_{n-1}})$. One concludes by observing that $\tau_{1,2}^{\alpha'_1} \dots \tau_{n-1,n}^{\alpha'_{n-1}} = \tau_{1,2} \dots \tau_{n-1,n}$ forces $\alpha'_1 = \alpha'_2 = \dots = \alpha'_{n-1} = 1$. The claim about cardinality is not difficult, once it is observed that an element of $\text{Red}_R(c)$ is uniquely determined by: 1) the position of the long reflection $\rho_{i,\zeta}$, 2) the integer i , 3) a reduced R -decomposition of the Coxeter element $\bar{c}$.

There remains a finite number of exceptional types which are treated by computer. $\square$

Until the end of this section, we assume that W is irreducible.

Lemma 7.6. *Let $y \in Y^{\text{gen}}$, with label $(s_1, \dots, s_n)$. Let $i \in \{1, \dots, n\}$. Then*

$$l_R(\pi(s_1 \dots s_i)) = i.$$

Proof. Since $y \in Y^{\text{gen}}$, each s_j is a braid reflection, mapped under π to a reflection $r_j \in R$. Thus $l_R(\pi(s_1 \dots s_i)) \leq i$ and $l_R(\pi(s_{i+1} \dots s_n)) \leq n - i$. Since $\pi(s_1 \dots s_i) \pi(s_{i+1} \dots s_n)$ is a Coxeter element (Lemma 7.2), it has length n . This forces both inequalities to be equalities. $\square$

Lemma 7.7. *The restriction of π to the set S of simple elements is injective.*

Proof. Let s and s' be simples such that $\pi(s) = \pi(s')$. By Corollary 6.20, we may find $y, y' \in Y^{\text{gen}}$, with $\text{lbl}(y) = (s_1, \dots, s_n)$ and $\text{lbl}(y') = (s'_1, \dots, s'_n)$, and $i, j \in \{0, \dots, n\}$ such that $s = s_1 \dots s_i$ and $s' = s'_1 \dots s'_j$. One can clearly assume that $LL(y) = LL(y')$. By Lemma 7.6, we must have $i = j$. Applying Proposition 7.5 to $w = \pi(s)$, we may find $\beta \in B_i$ such that $(\pi(s_1), \dots, \pi(s_i)) \cdot \beta = (\pi(s'_1), \dots, \pi(s'_i))$. Similarly, we find $\beta' \in B_{n-i}$ such that $(\pi(s_{i+1}), \dots, \pi(s_n)) \cdot \beta' = (\pi(s'_{i+1}), \dots, \pi(s'_n))$. View $B_i \times B_{n-i}$ as a subgroup of B_n (the first factor braids the first i strings, the second factors braids the $n - i$ last strings), and set $\beta'' := (\beta, \beta') \in B_n$. We have $\pi^n(\text{lbl}(y)) \cdot \beta'' = \pi^n(\text{lbl}(y'))$. Applying the theorem, this implies that $\text{lbl}(y) \cdot \beta'' = \text{lbl}(y')$. Clearly, β'' does not modify the product of the first i terms of the labels. Thus $s = s'$. $\square$

Definition 7.8. Let $x \in E_n$. Let $\underline{s} = (s_1, \dots, s_k)$ be a finite sequence of simple elements. We say that x and $\underline{s}$ are *compatible* if the following conditions are satisfied:

- The configuration x consists of k distinct points.
- Let $(x_1, \dots, x_k)$ be the ordered support of x . For all i , set

$$s'_i := \begin{cases} s_i & \text{if } i = n \text{ or } \text{re}(x_i) < \text{re}(x_{i+1}) \\ s_i s_{i+1}^{-1} & \text{if } i < n \text{ and } \text{re}(x_i) = \text{re}(x_{i+1}). \end{cases}$$

Then, for all i , s'_i is simple and $l(s'_i)$ is the multiplicity of x_i in x .

- We have $s'_1 \dots s'_k = \delta$.

In the following theorem, S^* denotes the set of finite sequences of elements of S .

Theorem 7.9. *The map $Y \rightarrow E_n \times S^*, y \mapsto (LL(y), \text{lbl}(y))$ is injective. Its image is the set of compatible pairs $(x, \underline{s}) \in E_n \times S^*$.*

Proof. (1) Injectivity.

The injectivity on Y^{gen} follows directly from Corollary 6.19 and Theorem 7.4.

Let $y, y' \in Y$ with $LL(y) = LL(y')$ and $lbl(y) = lbl(y')$. We have to prove that $y = y'$. Let $(x_1, \dots, x_k)$ and $(n_1, \dots, n_k)$ be the ordered support and multiplicities of $LL(y)$. Let $(t_1, \dots, t_k) := lbl(y)$.

For simplicity, we begin with the case when $\{x_1, \dots, x_k\} \in E_k^{\text{gen}}$, i.e., we assume that $\text{re}(x_1) < \text{re}(x_2) < \dots < \text{re}(x_k)$. Let $(y_m)_{m \geq 0}$ be sequences of points in Y^{gen} converging to y . Let $(s_1^{(m)}, \dots, s_n^{(m)}) := lbl(y_m)$. For m large enough, Hurwitz rule implies that $s_1^{(m)} \dots s_{n_1}^{(m)} = t_1$, $s_{n_1+1}^{(m)} \dots s_{n_1+n_2}^{(m)} = t_2$, etc... Similarly, choose a sequence (y'_m) of points in Y^{gen} converging to y' ; one has the corresponding properties for $(s_1^{(m)'}, \dots, s_n^{(m)'}) := lbl(y'_m)$. For m large, $(s_1^{(m)}, \dots, s_{n_1}^{(m)})$ and $(s_1^{(m)'}, \dots, s_{n_1}^{(m)'})$ are two reduced R -decompositions of the simple t_1 . By Proposition 7.5, they are in the same B_{n_1} Hurwitz orbit. Similarly, $(s_{n_1+1}^{(m)}, \dots, s_{n_1+n_2}^{(m)})$ and $(s_{n_1+1}^{(m)'}, \dots, s_{n_1+n_2}^{(m)'})$ are in the same B_{n_2} Hurwitz orbit, etc... We have proved that, for m large enough, $lbl(y_m)$ and $lbl(y'_m)$ are Hurwitz transformed by a braid β_m which lies in the natural subgroup $B_{n_1} \times B_{n_2} \times \dots \times B_{n_k}$ of B_n . One concludes noting that this subgroup belongs to the ramification group for the B_n -action on y .

The case when $\{x_1, \dots, x_k\} \notin E_k^{\text{gen}}$ may be reduced to the previous case by some suitable perturbation.

(2) Image.

Let $y \in Y$ be such that $LL(y) = x$ and $lbl(y) = \underline{s}$. Let ε be a small positive real number. We associate to $\underline{s} = s_1, \dots, s_k$ a sequence as in Definition 7.8. Clearly, $(s'_1, \dots, s'_k)$ is the label of $e^{-\sqrt{-1}\varepsilon}y$. Thus the s'_i are simple and, by Corollary 6.17, their product is δ . The statement about $l(s'_i)$ follows from Corollary 6.16.

Conversely, given any compatible x and $\underline{s} := (s_1, \dots, s_k)$, let us consider x' generic close to x and $y' \in LL^{-1}(x')$. For each i , choose a reduced R -decomposition of $\pi(s_i)$ and concatenate these decomposition to obtain a R -decomposition of c and, via Theorem 7.4, a possible label $\underline{s}'$ for a point in $LL^{-1}(x')$ (i.e., a point y'' in the Hurwitz orbit of y'). To conclude, degenerate y'' to a point with y with $LL(y) = x$ and apply Hurwitz rule. $\square$

Theorem 7.9 says that Y is in one-to-one correspondence with the set of compatible pairs $(x, \underline{s})$. Because of Theorem 7.4, we may actually replace $\underline{s}$ by the corresponding factorisation of c in W . We now have all the necessary tools to effectively work in Y . A final minor lemma will be needed:

Lemma 7.10. *Let $y \in Y$. Let $T = (y, z, L)$ and $T' = (y, z', L')$ be two tunnels in L_y such that $b_T = b_{T'}$. Then T and T' cross the same intervals among $I_1, \dots, I_n$ (in other words, T and T' are homotopic as tunnels drawn in L_y).*

Proof. Up to perturbing y , we may assume that $y \in Y^{\text{gen}}$. Let $(s_1, \dots, s_n) := lbl(y)$, and $(r_1, \dots, r_n) := \pi^n(lbl(y))$. Let $i_1, \dots, i_l$ (resp. $j_1, \dots, j_m$) be the successive indices of the intervals among $I_1, \dots, I_n$ crossed by T (resp. T'). We have $b_T = s_{i_1} \dots s_{i_l}$ and $b_{T'} = s_{j_1} \dots s_{j_m}$. Assuming that $b_T = b_{T'}$, we obtain $s_{i_1} \dots s_{i_l} = s_{j_1} \dots s_{j_m}$ and $r_{i_1} \dots r_{i_l} = r_{j_1} \dots r_{j_m}$. Let $w := r_{i_1} \dots r_{i_l}$. By Lemma 7.3, $l = m$ and we have

$$(*) \quad \ker(w - 1) = \bigcap_{k=1}^l \ker(r_{i_k} - 1) = \bigcap_{k=1}^l \ker(r_{j_k} - 1).$$

Assume that $(i_1, \dots, i_l) \neq (j_1, \dots, j_l)$. We may find $j \in \{j_1, \dots, j_l\}$ such that, for example, $i_1 < \dots < i_k < j < i_{k+1} < \dots < i_l$. Noting that the element $r_{i_1} \dots r_{i_k} r_j r_{i_{k+1}} \dots r_{i_l}$ is a parabolic Coxeter element, we deduce that $\ker(r_j - 1) \not\subseteq \ker(w - 1)$. This contradicts $(*)$. $\square$

8. THE DUAL BRAID MONOID

Here again, W is an irreducible well-generated complex reflection groups. We keep the notations from the previous sections. Recall that an element $b \in B$ is *simple* if $b = b_T$ for some tunnel T , and that the set of simple elements is denoted by S .

Definition 8.1. The *dual braid monoid* is the submonoid M of B generated by S .

Consider the binary relation $\preceq$ defined on S as follows. Let s and s' be simple elements. We write $s \preceq s'$ if and only if there exists $(z, y) \in W \setminus V^{\text{reg}}$, $L, L' \in \mathbb{R}_{\geq 0}$ with $L \leq L'$ such that (y, z, L) is a tunnel representing s and (y, z, L') is a tunnel representing s' .

This section is devoted to the proof of:

Theorem 8.2. *The monoid M is a Garside monoid, with set of simples S and Garside element δ . The relation $\preceq$ defined above on S is the restriction to S of the left divisibility order in M .*

The monoid M generates B , which inherits a structure of Garside group.

Remark 8.3. The theorem blends two results of different nature: one is about the Garside structure of a certain monoid $\mathbf{M}(P_c)$ (see Lemma 8.11 for a presentation of $\mathbf{M}(P_c)$); the other one identifies M with $\mathbf{M}(P_c)$. The latter essentially amounts to writing a presentation for B . It is a substitute for Brieskorn's presentation theorem, [15], except that our presentation involves *dual braid relations* instead of Artin-Tits braid relations.

Several results mentioned in the introduction follow from this theorem:

Theorem 0.4 does not assume irreducibility, but follows immediately from the irreducible case since direct products of Garside groups are Garside groups.

Another important consequence of Theorem 8.2 is that one may associate a practical contractible simplicial complex acted on by B . To explain this, we need some additional terminology, imitating [10] and [21]:

Definition 8.4. Let G be a group, let Σ be a subset of G . We denote by $\text{Flag}(G, \Sigma)$ the simplicial complex as follows:

- (0) the set of vertices is G ,
- (1) the 1-simplices are the pairs $\{g, h\}$ of distinct elements of G such that $g^{-1}h \in \Sigma$ or $h^{-1}g \in \Sigma$,
- (k) for all $k > 1$, a $(k+1)$ -tuple $\{g_0, \dots, g_k\}$ of distinct elements of G is a k -simplex if and only if $\{g_i, g_j\}$ is a 1-simplex for all $i \neq j$.

In other words, $\text{Flag}(G, \Sigma)$ is the only flag complex whose 1-skeleton is the Cayley graph of (G, Σ) .

The following result is due, in this formulation, to Charney-Meier-Whittlesey. It is inspired by the work of Bestvina, [10], on spherical type Artin monoid (which itself is based on the work of Deligne). It was also implicitly used in several articles by T. Brady (e.g. [12]).

Theorem 8.5 ([21]). *Let G be a Garside group with set of simples Σ . Then $\text{Flag}(G, \Sigma)$ is contractible.*

Corollary 8.6. *The simplicial complex $\text{Flag}(B, S)$ is contractible.*

The strategy of proof of Theorem 8.2 is very similar to the one in [6].

Proposition 8.7. *For all $s \in S$, we have $l(s) = l_R(\pi(s))$. For all $s, s' \in S$, the following statements are equivalent:*

- (i) $s \preceq s'$,
- (ii) $\exists s'' \in S, ss'' = s'$,

(iii) $\pi(s) \preceq_R \pi(s')$.

Proof. The first statement follows from Lemma 7.6.

(i) $\Rightarrow$ (ii). Assume that $s \preceq s'$ and choose $(x, y) \in W \setminus V^{\text{reg}}$ and $L \leq L'$ such that (y, x, L) is a tunnel representing s and (y, x, L') is a tunnel representing s' . Then $(y, x + L, L' - L)$ is a tunnel representing $s'' \in S$ such that $ss'' = s'$.

(ii) $\Rightarrow$ (iii). The natural length function l is additive on B . Thus, under (ii), we have $l(s) + l(s'') = l(s')$. On the other hand, for all $\sigma \in S$, $l(\sigma) = l_R(\pi(\sigma))$. Thus $l_R(\pi(s)) + l_R(\pi(s'')) = l_R(\pi(s'))$. Since $\pi(s'') = \pi(s)^{-1}\pi(s')$, this implies that $\pi(s) \preceq_R \pi(s')$.

(iii) $\Rightarrow$ (i). We may find $y, y' \in Y^{\text{gen}}$, with $lbl(y) = (s_1, \dots, s_n)$ and $lbl(y') = (s'_1, \dots, s'_n)$ such that $s = s_1 s_2 \dots s_{l(s)}$ and $s' = s'_1 s'_2 \dots s'_{l(s')}$. Set $w := \pi(s)$, $w' := \pi(s')$ and, for $i = 1, \dots, n$, set $r_i := \pi(s_i)$ and $r'_i := \pi(s'_i)$. Assuming (iii), we may find $r''_1, \dots, r''_{l(s')-l(s)} \in R$ such that $r_1 \dots r_{l(s)} r''_1 \dots r''_{l(s')-l(s)} = r'_1 \dots r'_{l(s')}$. The sequences

$$(r_1, \dots, r_{l(s)}, r''_1, \dots, r''_{l(s')-l(s)})$$

and

$$(r'_1, \dots, r'_{l(s')})$$

both lie in $\text{Red}_R(w')$. Since $w' \preceq c$, both sequences lie in the same Hurwitz orbit (Proposition 7.5). Thus

$$(r'_1, \dots, r'_n)$$

and

$$(r_1, \dots, r_{l(s)}, r''_1, \dots, r''_{l(s')-l(s)}, r'_{l(s')+1}, \dots, r'_n)$$

are transformed one onto the other by Hurwitz action of a braid $\beta \in B_n$ only braiding the first $l(s')$ strands. The Hurwitz transform of

$$lbl(y') = (s'_1, \dots, s'_n)$$

by β is the label

$$(s''_1, \dots, s''_n)$$

of some $y'' \in Y^{\text{gen}}$. Since the braid only involves the first $l(s')$ strands, $s''_1 \dots s''_{l(s')} = s'_1 \dots s'_{l(s')} = s'$. One has $\pi(s''_i) = r_i$ for $i = 1, \dots, l(s)$, thus $\pi(s''_1 \dots s''_{l(s)}) = \pi(s_1 \dots s_{l(s)})$. By Lemma 7.7, this implies $s''_1 \dots s''_{l(s)} = s_1 \dots s_{l(s)} = s$. For x with small enough real and imaginary parts, one may find real numbers L, L' with $0 < L \leq L'$ such that (y'', x, L) is a tunnel with associated simple $s''_1 \dots s''_{l(s)} = s$ and (y'', x, L') is a tunnel with associated simple $s''_1 \dots s''_{l(s')} = s'$. $\square$

Proposition 8.8. *The map π restricts to an isomorphism $(S, \preceq) \xrightarrow{\sim} ([1, c], \preceq_R)$. In particular, $\preceq$ is an order relation on S .*

Proof. The previous proposition, applied to $s' = \delta$, proves that $\pi(S) \subseteq [1, c]$. It also proves that π induces a morphism of sets with binary relations $(S, \preceq) \rightarrow ([1, c], \preceq_R)$. The injectivity is Lemma 7.7.

Surjectivity: Choose $y \in Y^{\text{gen}}$. Let $(s_1, \dots, s_n) := lbl(y)$. Let $r_i := \pi(s_i)$. We have $(r_1, \dots, r_n) \in \text{Red}_R(c)$. Let $w \in [1, c]$. We may find $(r'_1, \dots, r'_n) \in \text{Red}_R(c)$ such that $r'_1 \dots r'_{l_T(w)} = w$. By Proposition 7.5, $(r'_1, \dots, r'_n)$ is Hurwitz transformed of $(r_1, \dots, r_n)$, thus there exists $y' \in Y^{\text{gen}}$ such that $\pi_*(lbl(y')) = (r'_1, \dots, r'_n)$. The simple element which is the product of the first $l(w)$ terms of $lbl(y')$ is in $\pi^{-1}(w)$. $\square$

We have the key lemma:

Lemma 8.9. *The poset $([1, c], \preceq_R)$ is a lattice.*

Proof. The real case is done in [3] – a recent beautiful case-free argument has been found by Brady-Watt, [14]. The case of $G(e, e, r)$ is done in [6]. The remaining cases have been checked by computer. $\square$

Definition 8.10. The *dual braid relations* with respect to W and c are all the formal relations of the form

$$rr' = r'r'',$$

where $r, r', r'' \in R$ are such that $r \neq r', rr' \in [1, c]$, and the relation $rr' = r'r''$ holds in W .

Clearly, dual braid relations only involve reflections in $R \cap [1, c]$. When W is complexified real, $R \subseteq [1, c]$. This does not hold in general (see the tables at the end of the article).

As in [3] (see also [5, Section 1]), one endows $[1, c]$ with a partial product and obtains a monoid $\mathbf{M}(P_c)$. From Lemma 8.9, we will deduce that $\mathbf{M}(P_c)$ is a Garside monoid. We will also identify $\mathbf{M}(P_c)$ with M . The following lemma generalises [3, Theorem 2.1.4]:

Lemma 8.11. *Let $R_c := R \cap [1, c]$. The monoid $\mathbf{M}(P_c)$ admits the monoid presentation*

$$\mathbf{M}(P_c) \simeq \langle R_c \mid \text{dual braid relations} \rangle.$$

Remark 8.12. When viewed as a group presentation, the presentation of the lemma is a presentation for $\mathbf{G}(P_c)$. As soon as we prove Theorem 8.2, Lemma 8.11 will give an explicit presentation for B . A way to reprove Theorem 0.6 is by simplifying the (redundant) presentation given by the lemma. This does not involve any monodromy computation.

Proof of Lemma 8.11. By definition, $\mathbf{M}(P_c)$ admits the presentation with generators R_c and a relation $r_1 \dots r_k = r'_1 \dots r'_k$ for each pair $(r_1, \dots, r_k), (r'_1, \dots, r'_k)$ of reduced R -decompositions of the same element $w \in [1, c]$. Call these relations *Hurwitz relations*. By transitivity of the Hurwitz action on $\text{Red}_R(c)$ (Proposition 7.5), the Hurwitz relations are consequences of the dual braid relations. The dual braid relations clearly hold in $\mathbf{M}(P_c)$ (to see this, complete (r, r') to an element $(r, r', r_3, \dots, r_n) \in \text{Red}_R(c)$). This proves the lemma. $\square$

Proof of Theorem 8.2. Set $R_c := R \cap [1, c]$. We are in the situation of subsection 0.4 in [3], (W, R_c) is a *generated group* and c is *balanced* (one first observes that $\{w \mid w \preceq_{R_c} c\} = \{w \mid w \preceq_R c\}$ and $\{w \mid c \succ_{R_c} w\} = \{w \mid c \succ_R w\}$; one concludes noting that c is balanced with respect to (W, R) , which is immediate since R is invariant by conjugacy). By Lemma 8.9 and [3, Theorem 0.5.2], the premonoid $P_c := [1, c]$ (together with the natural partial product) is a Garside premonoid. We obtain a Garside monoid $\mathbf{M}(P_c)$.

By Proposition 8.8, the restriction of π is a bijection from S to P_c . Let ϕ be the inverse bijection. Let $w, w' \in P_c$. Assume that the product $w.w'$ is defined in P_c . Let w'' be the value of this product. One has $w \preceq_R w''$, thus $\phi(w) \preceq \phi(w'')$ (using again Proposition 8.8), and we may find b' in S such that $\phi(w)b' = \phi(w'')$ (Proposition 8.7). Claim: $b' = \phi(w')$. Indeed, b' and $\phi(w')$ are two simple elements whose image by π is $w^{-1}w''$; one concludes using Lemma 7.7. This proves that ϕ induces a premonoid morphism $P_c \rightarrow S$ (where S is equipped with the restriction of the monoid structure), thus induces a monoid morphism $\mathbf{M}(P_c) \rightarrow M$ and a group morphism $\Phi : \mathbf{G}(P_c) \rightarrow B$.

Let us prove that Φ is an isomorphism. Choose a basepoint $y \in Y^{\text{gen}}$. Let $\gamma_1, \dots, \gamma_n$ be generators of $\pi_1(Y - \mathcal{K}, y)$. Let $(s_1, \dots, s_n)$ be the label of y . Let us reinterpret the presentation from Theorem 4.4 in terms of Hurwitz action. Since $LL : Y - \mathcal{K} \rightarrow E_n^{\text{gen}}$ is a covering, $\pi_1(Y - \mathcal{K}, y)$ may be identified with a subgroup $H \subseteq B_n$. The generators $f_1, \dots, f_n$ in Theorem 4.4 may be chosen to be $s_1, \dots, s_n$, and the monodromy automorphism $\phi_1, \dots, \phi_m$ are obtained by Hurwitz action on $s_1, \dots, s_n$. Let $h \in H$. Let $(s'_1, \dots, s'_n) := h(s_1, \dots, s_n)$ (by this, we mean the Hurwitz action of h on the free group generated by $s_1, \dots, s_n$; the s'_i 's are words in the s_i 's). Call *Van*

Kampen relations the relations of the type $s'_i = s_i$, for any $i \in \{1, \dots, n\}$, $h \in H$, and s'_i obtained as above. We have

$$B \simeq \langle s_1, \dots, s_n \mid \text{Van Kampen relations} \rangle.$$

The map π induces a bijection from $A := \pi^{-1}(R_c) \subseteq S$ to R_c . Let $r_1, \dots, r_n$ be the images of $s_1, \dots, s_n$. By transitivity of Hurwitz action on $\text{Red}_R(c)$, the group $\mathbf{G}(P_c)$ is generated by $r_1, \dots, r_n$, the remaining generators in the presentation of Lemma 8.11 appearing as conjugates of $r_1, \dots, r_n$ (by successive use of dual braid relations). Our generating sets are compatible and the morphism

$$\Phi : \mathbf{G}(P_c) \simeq \langle R_c \mid \text{dual braid relations} \rangle \rightarrow B \simeq \langle s_1, \dots, s_n \mid \text{Van Kampen relations} \rangle$$

is defined by $r_i \mapsto s_i$. Add to the presentation of B formal generators indexed by

$$A - \{s_1, \dots, s_n\}$$

as well as the dual braid relations $\pi(r)\pi(r') = \pi(r')\pi(r'')$. Since the relations already hold in $\mathbf{G}(P_c)$, they hold in B , and we obtain a new presentation

$$B \simeq \langle A \mid \text{Van Kampen relations on } \{s_1, \dots, s_n\}, \text{ dual braid relations on } A \rangle.$$

To conclude that Φ is an isomorphism, it is enough to observe that the dual braid relations encode the full Hurwitz action of B_n of $(s_1, \dots, s_n)$, while the Van Kampen relations encode the action of $H \subseteq B_n$: thus Van Kampen relations are consequences of dual braid relations, and $\mathbf{G}(P_c)$ and B are given by equivalent presentations.

Since Φ is an isomorphism and $\mathbf{M}(P_c)$ naturally embeds in $\mathbf{G}(P_c)$ (this is a crucial property of Garside monoids, [23]), $\mathbf{M}(P_c)$ is isomorphic to its image M in B . The rest of theorem is clear. $\square$

As mentioned earlier, one may view B as a “reflection group”, generated by the set $\mathcal{R}$ of all braid reflections. An element of M is in $\mathcal{R}$ if and only if it has length 1 for the natural length function, or equivalently if it is an *atom* (*i.e.*, an element which has no strict divisor in M except the unit). By Proposition 7.5 and Theorem 7.4, there is a bijection between $\text{Red}_R(c)$ and the image of $lbl : Y^{\text{gen}} \rightarrow \mathcal{R}^n$. This image is clearly contained in $\text{Red}_{\mathcal{R}}(\delta)$. The conjecture below claims an analogue in B of the transitivity of the Hurwitz action on $\text{Red}_R(c)$ (δ is the natural substitute for a Coxeter element in B). It implies that any element in $\text{Red}_{\mathcal{R}}(\delta)$ is the label of some $y \in Y^{\text{gen}}$.

Conjecture 8.13. *The Hurwitz action of B_n on $\text{Red}_{\mathcal{R}}(\delta)$ is transitive.*

9. CHAINS OF SIMPLES

Here again, W is an irreducible well-generated complex reflection group, and the notations from the previous section are still in use. For $k = 0, \dots, n$, we denote by $\mathcal{C}_k$ the set of (strict) chains in $S - \{1\}$ of cardinal k , *i.e.*, the set of k -tuples $(c_1, \dots, c_k)$ in S^k such that

$$1 \prec c_1 \prec \dots \prec c_k,$$

or equivalently the set of k -tuples $(c_1, \dots, c_k)$ in M^k such that


$$1 \prec c_1 \prec \dots \prec c_k \preceq \delta.$$

It is convenient to write $\{1 \prec c_1 \prec \dots \prec c_k\}$ instead of $(c_1, \dots, c_k)$.

We set $\mathcal{C} := \bigsqcup_{k=0}^n \mathcal{C}_k$.

Let $C := \{1 \prec c_1 \prec \dots \prec c_k\} \in \mathcal{C}$. We say that $y \in Y$ *represents* C if there exists $x \in U_y$ and real numbers $L_1, \dots, L_k$ such that $0 < L_1 < \dots < L_k$ and, for $i = 1, \dots, k$, (y, x, L_i) is a tunnel representing C_i .

Example: if $LL(y)$ is as in the illustration below and $(s_1, \dots, s_5) = lbl(y)$, y represents $1 \prec s_1 \prec s_1 s_3 \prec s_1 s_3 s_4 \prec s_1 s_3 s_4 s_5$, $1 \prec s_2 \prec s_2 s_4 \prec s_2 s_4 s_5$, $1 \prec s_2 \prec s_2 s_5$ and their subchains, but does not represent $1 \prec s_2 \prec s_2 s_3$ nor $1 \prec s_3 \prec s_3 s_5$.



Definition 9.1. For all $C \in \mathcal{C}$, we set $Y_C := \{y \in Y \mid y \text{ represents } C\}$.

To illustrate this notion, we observe that implication (ii) $\Rightarrow$ (i) from Proposition 8.7 expresses that for all $C \in \mathcal{C}_2$, Y_C is non-empty. Based on the results from the previous sections, this easily generalises to:

Lemma 9.2. For all $C \in \mathcal{C}$, the space Y_C is non-empty.

The goal of this section is to prove:

Proposition 9.3. For all $C \in \mathcal{C}$, the space Y_C is contractible.

This technical result will be used in Section 10, when studying the nerve of an open covering of the universal cover of $W \setminus V^{\text{reg}}$: we will need to prove that certain non-empty intersections of open sets are contractible, and these intersections will appear as fiber bundles over some Y_C , with contractible fibers.

The proposition is not very deep nor difficult but somehow *inconvenient* to prove since the retraction will be described via LL , through ramification points. The following particular cases are easier to obtain:

- If C is the chain $1 \prec \delta$, then $Y_C = Y \simeq \mathbb{C}^{n-1}$.
- More significantly, let W be a complex reflection group of type A_2 . Up to renormalisation, the discriminant is $X_2^2 + X_1^3$. Identify Y with $\mathbb{C}$. For all $y \in \mathbb{C}$, $LL(y) = \{\pm(-y)^{3/2}\}$. In particular, $LL(1) = \{\pm\sqrt{-1}\}$. Let s be the simple element represented by the tunnel with $y = 1$, $x = -1$ and $L = 2$. Let C be the chain $1 \prec s$. Then Y_C is the open cone consisting of non-zero elements of $\mathbb{C}$ with argument in the open interval $(-\pi/3, \pi/3)$.
- Assume that $C \in \mathcal{C}_n$. All points in Y_C are generic. Consider the map $Y^{\text{gen}} \rightarrow \mathcal{C}_n \times E_n$ sending y to the pair $(\{1 \prec s_1 \prec s_1 s_2 \prec \dots \prec s_1 s_2 \dots s_n = \delta\}, LL(y))$, where $(s_1, \dots, s_n) = lbl(y)$. This map is an homeomorphism. The $(Y_C)_{C \in \mathcal{C}_n}$ are the connected components of Y^{gen} . Each of these components is homeomorphic to E_n , which is contractible. These $(Y_C)_{C \in \mathcal{C}_n}$ are some analogues of chambers.

Regular coverings satisfy a unique homotopy lifting property. The map $LL : Y \rightarrow E_n$ is not regular on $\mathcal{K}$ but is “stratified regular”, where E_n is endowed with the discriminant stratification

$$E_n = \bigsqcup_{\lambda \models n} S_\lambda$$

and Y with the corresponding stratification (see Corollary 5.8 and the subsequent discussion.) A consequence of this is that paths with non-decreasing ramification admit unique lifts:

Lemma 9.4. Let $\gamma : [0, 1] \rightarrow E_n$ be a path. For all $t \in [0, 1]$, denote by S_t the stratum of E_n containing $\gamma(t)$. Assume that $\forall t, t' \in [0, 1], t \leq t' \Rightarrow \overline{S_t} \supseteq S_{t'}$. Let $y_0 \in LL^{-1}(\gamma(0))$. There exists a unique path $\tilde{\gamma} : [0, 1] \rightarrow Y$ such that $\gamma = LL \circ \tilde{\gamma}$ and $\tilde{\gamma}(0) = y_0$.

Proof. Let $0 = t_0 < t_1 < \dots < t_m = 1$ be such that S_t is constant on each $[t_{i-1}, t_i)$. By an obvious induction on m , it suffices to show that there exists a unique $\tilde{\gamma} : [0, t_1] \rightarrow Y$ such that,

for all $t \in [0, t_1)$, $\gamma(t) = LL \circ \tilde{\gamma}(t)$ and $\tilde{\gamma}(0) = y_0$. Because ramification is constant over $[0, t_1)$, there is a unique continuous lift over $[0, t_1)$. Hurwitz rule determines the label of the unique $\gamma(t_1)$ that makes the lift continuous over $[0, t_1]$. $\square$

In the following proposition, if $A \subseteq LL(y)$ is a submultiset, the *deep label* of A is the sequence $(t_1, \dots, t_p)$ of labels (with respect to y) of points in the support of A which are deep with respect to A (since these points may not be deep in $LL(y)$, the deep label of A is not necessarily a subsequence of the deep label of y).

Proposition 9.5. *Let $C = \{1 \prec c_1 \prec c_2 \prec \dots \prec c_m\}$ be a chain in $\mathcal{C}_m$. Let $y \in Y$, let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$ and $(s_1, \dots, s_k)$ be the label of y . The following assertions are equivalent:*

- (i) $y \in Y_C$.
- (ii) *There exists a partition of $LL(y)$ into $m+1$ submultisets $A_0, \dots, A_m$ such that:*
 - (a) *For all $i, j \in \{1, \dots, m\}$ with $i < j$, for all $x \in A_i$ and all $x' \in A_j$, we have $\text{re}(x) < \text{re}(x')$.*
 - (b) *For all $x \in A_0$, one has*

$$\text{re}(x) < \min_{x' \in A_1 \cup \dots \cup A_m} \text{re}(x')$$

or

$$\text{im}(x) < \min_{x' \in A_1 \cup \dots \cup A_m} \text{im}(x')$$

or

$$\text{re}(x) > \max_{x' \in A_1 \cup \dots \cup A_m} \text{re}(x').$$


- (c) *For all $i \in \{1, \dots, m\}$, the product of the deep label of A_i is c_i .*

Moreover, in situation (ii), the partition $LL(y) = A_0 \sqcup \dots \sqcup A_m$ is uniquely determined by y and C .

The picture below illustrates the proposition for particular y and C . Here $\text{lbl}(y) = (s_1, \dots, s_5)$ and the considered chain is

$$C = \{1 \prec s_2 \prec s_2 s_4\}.$$

We have chosen tunnels $T_1 = (y, x, L_1)$ and $T_2 = (y, x, L_2)$, with $L_1 < L_2$, such that $b_{T_1} = s_2$ and $b_{T_2} = s_2 s_4$. The dotted lines represent these tunnels, as well as the vertical half-lines above x , $x + L_1$ and $x + L_2$. They partition the complex line into three connected components; the partition $A_0 \sqcup A_1 \sqcup A_2$ is the associated partition of $LL(y)$. It is clear the possibility of drawing the tunnels is subject precisely to the conditions on $A_0 \sqcup A_1 \sqcup A_2$ expressed in the proposition.



Proof. (i) $\Rightarrow$ (ii): $(y, z, L_1), \dots, (y, z, L_m)$ be tunnels representing the successive non-trivial terms of $C = \{1 \prec c_1 \prec \dots \prec c_m\}$. Set $z_0 := z$ and, for $i = 1, \dots, m$, $z_i := z + L_i$. We have

$$\text{re}(z_0) < \text{re}(z_1) < \dots < \text{re}(z_m)$$

and

$$\text{im}(z_0) = \text{im}(z_1) = \dots = \text{im}(z_m).$$

For $i \in \{1, \dots, m\}$, let A_i be the submultiset of $LL(y)$ consisting of points x such that $\text{re}(z_{i-1}) < \text{re}(x) < \text{re}(z_i)$ and $\text{im}(x) > \text{im}(z_0)$. Let A_0 be the complement in $LL(y)$ of $A_1 \cup \dots \cup A_m$. One easily checks (a), (b) and (c).

(ii) $\Rightarrow$ (i): Conversely, assume we are given a partition $A_0 \sqcup A_1 \sqcup \dots \sqcup A_m$ satisfying conditions (a) and (b). One may recover tunnels $(y, z, L_1), \dots, (y, z, L_m)$ such that the above construction yields the partition $A_0 \sqcup A_1 \sqcup \dots \sqcup A_m$. Condition (c) then implies that the tunnels represent the elements of C , thus that $y \in Y_C$.

Unicity of the partition: this follows from condition (c) and Lemma 7.10. $\square$

Lemma 9.6. *Let $C = \{1 \prec c_1 \prec c_2 \prec \dots \prec c_m\} \in \mathcal{C}$. Define Y_C^0 as the subspace of Y_C consisting of points y whose associated partition $A_0, \dots, A_m$ (from Proposition 9.5 (ii)) satisfies the following conditions:*


- for $i = 0, \dots, m$, the support of A_i is a singleton $\{a_i\}$ and,
- $\text{re}(a_0) = \min_{i=1, \dots, m} \text{re}(a_i) - 1$ and $\text{im}(a_0) = \min_{i=1, \dots, m} \text{im}(a_i) - 1$.

Then Y_C^0 is contractible.

Proof. Let $y \in Y_C^0$. The support $(x_0, \dots, x_m)$ of $LL(y)$ is generic, thus the label $(s_0, \dots, s_m)$ of y coincides with the deep label, and we have $s_0 \dots s_m = \delta$ (Corollary 6.17). By Proposition 9.5, Condition (ii)(c), we have, for $i = 1, \dots, m$, $s_i = c_i$. Thus $s_0 = \delta(c_1 \dots c_m)^{-1}$. We have proved that the label of any $y \in Y_C^0$ must be $(\delta(c_1 \dots c_m)^{-1}, c_1, \dots, c_m)$. A consequence of Theorem 7.9 is that the map $Y_C^0 \rightarrow E_m^{\text{gen}}$ sending y to $(x_1, \dots, x_m)$ is an homeomorphism. One concludes with Lemma 3.2. $\square$

We may now proceed to the proof of Proposition 9.3. Let $C = \{1 \prec c_1 \prec \dots \prec c_m\} \in \mathcal{C}$. Let $y \in Y_C$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$, and $(n_1, \dots, n_k)$ be the multiplicities. Let $A_0, \dots, A_m$ be the partition of $LL(y)$ described in Proposition 9.5 (ii).

The picture below gives an idea of the retraction of Y_C onto Y_C^0 that will be explicitly constructed. It illustrates the motion of a given point $y \in Y_C$; the black dots indicate the support of $LL(y)$ and the arrows how this support moves during the retraction.



For $i = 1, \dots, m$, consider the multiset mass centre

$$a_i := \frac{\sum_{x \in A_i} x}{|A_i|}$$

(in this expression, A_i is viewed as a multiset: each x_j in A_i is taken n_j times, and $|A_i|$ is the multiset cardinal, i.e., the sum of the n_j such that $x_j \in A_i$).

For each $t \in [0, 1]$, let

$$\gamma_y(t) := A_0 \cup \bigcup_{i=1}^m \{(1-t)x + ta_i \mid x \in A_i\}$$

(here again, we consider the multiset union; in particular, the multicardinal of $\gamma_y(t)$ is constant, equal to n - i.e., $\gamma_y(t) \in E_n$). This defines a path in E_n .

The path γ_y satisfies the hypothesis of Lemma 9.4 and uniquely lifts to a path $\tilde{\gamma}_y$ in Y such that $\tilde{\gamma}_y(0) = y$. An easy consequence of Proposition 9.5 is that $\tilde{\gamma}_y$ is actually drawn in Y_C .

Let $y' := \tilde{\gamma}_y(1)$. Let

$$R := \min_{i=1,\dots,m} \operatorname{re}(a_i) = \operatorname{re}(a_1)$$

and

$$I := \min_{i=1,\dots,m} \operatorname{im}(a_i).$$

For all $x \in A_0$, we have $\operatorname{re}(x) < R$ or $\operatorname{im}(x) < I$ or $\operatorname{re}(x) > \operatorname{re}(a_m)$; denote by x' the complex number with the same real part as x and imaginary part $I - 1$; let $a_0 := R - 1 + \sqrt{-1}(I - 1)$. Consider the path $\beta_x : [0, 1] \rightarrow \mathbb{C}$ defined by:

$$\beta_x(t) := \begin{cases} (1 - 2t)x + 2tx' & \text{if } t \leq 1/2 \\ (2 - 2t)x' + (2t - 1)a_0 & \text{if } t \geq 1/2. \end{cases}$$

For all $t \in [0, 1]$, one has $\operatorname{re}(\beta_x(t)) < R$ or $\operatorname{im}(\beta_x(t)) < I$ or $\operatorname{re}(\beta_x(t)) > \operatorname{re}(a_m)$.

The path $\gamma'_y : [0, 1] \rightarrow E_n$ defined by

$$\gamma'_y(t) := \underbrace{\{a_1, \dots, a_1\}}_{|A_1| \text{ times}} \cup \dots \cup \underbrace{\{a_m, \dots, a_m\}}_{|A_m| \text{ times}} \cup \bigcup_{x \in A_0} \beta_x(t)$$

satisfies the hypothesis of Lemma 9.4 and lifts to a unique path $\tilde{\gamma}'_y$ in Y such that $\tilde{\gamma}'_y(0) = y'$. Once again, a direct application of Proposition 9.5 ensures that $\tilde{\gamma}'_y$ is actually in Y_C . The endpoint $y'' := \tilde{\gamma}'_y(1)$ lies in the subspace Y_C^0 of Corollary 9.6.

The map

$$\begin{aligned} \varphi : Y_C \times [0, 1] &\longrightarrow Y_C \\ (y, t) &\longmapsto (\tilde{\gamma}'_y \circ \tilde{\gamma}_y)(t) \end{aligned}$$

is a retraction of Y_C onto its contractible subspace Y_C^0 . Thus Y_C is contractible.

10. THE UNIVERSAL COVER OF $W \backslash V^{\text{reg}}$

The main result of this section is:

Theorem 10.1. *The universal cover of $W \backslash V^{\text{reg}}$ is homotopy equivalent to $\operatorname{Flag}(B, S)$.*

Combined with Corollary 8.6, this proves our main result Theorem 0.2 in the irreducible case. The reducible case follows.

To prove this theorem, we construct an open covering $\widehat{\mathcal{U}} = (\widehat{\mathcal{U}}_b)_{b \in B}$, such that the nerve of $\mathcal{U}$ is $\operatorname{Flag}(B, S)$ (Proposition 10.6) and intersections of elements of $\widehat{\mathcal{U}}$ are either empty or contractible (Proposition 10.7). Under these assumptions, a standard theorem from algebraic topology ([30, 4G.3]) gives the desired result.

As explained in the topological preliminaries, our “basepoint” $\mathcal{U}$ provides us with a model denoted $(\widetilde{W \backslash V^{\text{reg}}})_{\mathcal{U}}$ or simply $\widetilde{W \backslash V^{\text{reg}}}$ for the universal cover of $W \backslash V^{\text{reg}}$. Recall that, to any semitunnel T , one associates a path γ_T whose source is in $\mathcal{U}$, thus a point in $\widetilde{W \backslash V^{\text{reg}}}$. Moreover, we have a left action of $B = \pi_1(W \backslash V^{\text{reg}}, \mathcal{U})$ on $\widetilde{W \backslash V^{\text{reg}}}$. With these conventions, our open covering is very easy to define:

Definition 10.2. The set $\widehat{\mathcal{U}}_1$ is the subset of $\widetilde{W \backslash V^{\text{reg}}}$ of elements represented by semitunnels. For all $b \in B$, we set $\widehat{\mathcal{U}}_b := b\widehat{\mathcal{U}}_1$.

It is clear that two semitunnels represent the same point in $\widetilde{W \backslash V^{\text{reg}}}$ if and only if they are equivalent in the following sense:

Definition 10.3. Two semitunnels $T = (y, x, L)$ and $T' = (y', x', L')$ are *equivalent* if and only if $y = y'$, $x + L = x' + L'$ and the affine segment $[(x, y), (x', y)]$ is included in $\mathcal{U}$.

Let $T = (y, x, L)$ be a semitunnel. The point of $\widehat{\mathcal{U}}_1$ determined by T , or in other words the equivalence class of T , is uniquely determined by y , $x + L$ and

$$\lambda(T) := \inf \{ \lambda' \in [0, L] \mid [(x, y), (x + L - \lambda', y)] \subseteq \mathcal{U} \}.$$

The number λ is the infimum of the length of semitunnels in the equivalence class of T . This infimum is not a minimum (unless T is included in $\mathcal{U}$), since $(y, x + L - \lambda(T), \lambda(T))$ is not in general a semitunnel.

We consider the following subsets of $\widehat{\mathcal{U}}_1$:

- ($\mathcal{U}_1$) If $\lambda(T) = 0 = \min \{ \lambda' \in [0, L] \mid [x, x + L - \lambda'] \subseteq U_y \}$, then T is a tunnel, equivalent to $(y, x + L, 0)$. Elements of $\widehat{\mathcal{U}}_1$ represented by such tunnels of length 0 form an open subset denoted by $\mathcal{U}_1$. This subset is actually a sheet over $\mathcal{U}$ of the universal covering, corresponding to the trivial lift of the “basepoint”.
- ($\overline{\mathcal{U}}_1$) We denote by $\overline{\mathcal{U}}_1$ the subset consisting of points represented by tunnels with $\lambda(T) = 0$ (but without requiring that the “inf” is actually a “min”). We obviously have $\mathcal{U}_1 \subseteq \overline{\mathcal{U}}_1$, and $\overline{\mathcal{U}}_1$ is contained in the closure of $\mathcal{U}_1$.

Similarly, we set for all $b \in B$

$$\mathcal{U}_b := b\mathcal{U}_1, \quad \overline{\mathcal{U}}_b := b\overline{\mathcal{U}}_1.$$

Also, for any $y \in Y$, we denote by $\widehat{\mathcal{U}}_{b,y}$ (resp. $\mathcal{U}_{b,y}$, resp. $\overline{\mathcal{U}}_{b,y}$) the intersection of $\widehat{\mathcal{U}}_b$ (resp. $\mathcal{U}_b$, resp. $\overline{\mathcal{U}}_b$) with the fiber over y of the composed map $\widetilde{W \setminus V^{\text{reg}}} \rightarrow W \setminus V^{\text{reg}} \rightarrow Y$.

Lemma 10.4. *The family $(\overline{\mathcal{U}}_b)_{b \in B}$ is a partition of $\widetilde{W \setminus V^{\text{reg}}}$.*

Proof. It suffices to show that the projection $\overline{\mathcal{U}}_1 \rightarrow W \setminus V^{\text{reg}}$ is bijective. Any point $z \in W \setminus V^{\text{reg}}$ is the target of a unique equivalence class of semitunnels T with $\lambda(T) = 0$; depending on whether $z \in \mathcal{U}$ or not, the associated point will be in $\mathcal{U}_1$ or in $\overline{\mathcal{U}}_1 - \mathcal{U}_1$. $\square$

In particular, $(\widehat{\mathcal{U}}_b)_{b \in B}$ is a covering of $\widetilde{W \setminus V^{\text{reg}}}$.

Lemma 10.5. *For all $b \in B$, $\widehat{\mathcal{U}}_b$ is open and contractible.*

Proof. It is enough to deal with $b = 1$. That $\widehat{\mathcal{U}}_1$ is open is easy. Let $\mathcal{T}$ be the space of tunnels and $\sim$ the equivalence relation. As a set, $\widehat{\mathcal{U}}_1 \simeq \mathcal{T} / \sim$. Consider the map

$$\begin{aligned} \phi : \mathcal{T} \times [0, 1] &\longrightarrow \mathcal{T} \\ (T = (y, x, L), t) &\longmapsto (y, x, L - t\lambda(T)) \end{aligned}$$

If $T \sim T'$, then $\forall t, \phi(T, t) \sim \phi(T', t)$. Thus ϕ induces a map $\overline{\phi} : \widehat{\mathcal{U}}_1 \times [0, 1] \rightarrow \widehat{\mathcal{U}}_1$. This map is continuous (this follows from the fact that λ induces a continuous function on $\widehat{\mathcal{U}}_1$) and $\overline{\phi}(T, t) = T$ if $T \in \overline{\mathcal{U}}_1$ or if $t = 0$. We have proved that $\widehat{\mathcal{U}}_1$ retracts to $\overline{\mathcal{U}}_1$. We are left with having to prove that $\overline{\mathcal{U}}_1$ is contractible. Inside $\overline{\mathcal{U}}_1$ lies $\mathcal{U}_1$ which is contractible, since it is a standard lift of the contractible space $\mathcal{U}$.

We conclude by observing that $\mathcal{U}_1$ and $\overline{\mathcal{U}}_1$ are homotopy equivalent. There is probably a standard theorem from semialgebraic geometry applicable here, but I was unable to find a proper reference. Below is a “bare-hand” argument: it explains how $\overline{\mathcal{U}}_1$ may be “locally retracted” inside $\mathcal{U}_1$ (constructing a global retraction seems difficult).

Both spaces have homotopy type of CW -complexes and to prove homotopy equivalence it is enough to prove that any continuous map $f : S^k \rightarrow \overline{\mathcal{U}}_1$ (where S^k is a sphere) may be

homotoped to a map $S^k \rightarrow \mathcal{U}_1$. Assume that there is a tunnel $T = (y, z, L)$ such that $(T/\sim) \in f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1)$. We have $L > 0$. For any $\varepsilon > 0$, let $B_\varepsilon(y)$ be the open ball of radius ε in Y around y , let $B_\varepsilon(z)$ be the affine interval $(z - \sqrt{-1}\varepsilon, z + \sqrt{-1}\varepsilon)$. For ε small enough, there is a unique continuous function $L_\varepsilon : B_\varepsilon(y) \times B_\varepsilon(z) \rightarrow \mathbb{R}$ such that $L_\varepsilon(y, z) = L$ and for all $(y', z') \in B_\varepsilon(y) \times B_\varepsilon(z)$, $(y', z', L_\varepsilon(y', z'))$ represents a point in $\overline{\mathcal{U}}_1 - \mathcal{U}_1$. The “half-ball”

$$H_\varepsilon := \{(y', z', L') \in B_\varepsilon(y) \times B_\varepsilon(z) \times \mathbb{R} \mid 0 \leq L' \leq L_\varepsilon(y', z')\}$$

is a neighbourhood of $T/\sim$ in $\overline{\mathcal{U}}_1$. Working inside this neighbourhood, one may homotope f to f' such that $f'(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1) \subseteq f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1) - \{T/\sim\}$. Compacity of $f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1)$ guarantees that one can iterate this process a finite number of times to get rid of all $f(S^k) \cap (\overline{\mathcal{U}}_1 - \mathcal{U}_1)$. $\square$

Proposition 10.6. *The nerve of $(\widehat{\mathcal{U}}_b)_{b \in B}$ is $\text{Flag}(B, S)$.*

Proof. Let $b, b' \in B$ such that $\widehat{\mathcal{U}}_b \cap \widehat{\mathcal{U}}_{b'} \neq \emptyset$. Let $T = (y, x, L)$ and $T' = (y', x', L')$ be semitunnels, representing points z and z' in $\widehat{\mathcal{U}}_1$, such that $bz = b'z'$. The image of z (resp. z') in $W \setminus V^{\text{reg}}$ is $(x + L, y)$ (resp. $(x' + L', y')$). Thus $x + L = x' + L'$ and $y = y'$. Up to permuting b and b' , we may assume that $L \geq L'$. Since $x' = x + L - L'$ is in U_y , $T'' := (y, x, L - L')$ is a tunnel, representing a simple element b'' . The tunnel T is a concatenation of T'' and T' . This implies that $z = b''z'$ and $b'z' = bz = bb''z'$. By faithfulness of the B -action on the orbit of z , we conclude that $bb'' = b'$.

We have proved that the 1-skeletons of the nerve and of $\text{Flag}(B, S)$ coincide. To conclude, it remains to check that the nerve is a flag complex. Let $C \subseteq B$ be such that for all $b, b' \in C$, either $b^{-1}b'$ or $b'^{-1}b$ is simple. We have to prove that $\bigcup_{b \in C} \widehat{\mathcal{U}}_b \neq \emptyset$.

Let $c_0, \dots, c_m$ be the elements of C , numbered according to the total ordering on C induced by $\prec$:

$$c_0 \prec c_1 \prec c_2 \prec \dots \prec c_m \preccurlyeq c_0\delta.$$

Up to left-dividing each term by c_0 , we may assume that $c_0 = 1$. Let $y \in Y_C$. We may find $x, L_1, \dots, L_m$ such that (y, x, L_i) represents c_i for all i . The point of the universal cover represented by (y, x, L_m) belongs to $\bigcup_{b \in C} \widehat{\mathcal{U}}_b$. $\square$

Proposition 10.7. *Let C be a subset of B such that $\bigcap_{b \in C} \widehat{\mathcal{U}}_b \neq \emptyset$. Then $\bigcap_{b \in C} \widehat{\mathcal{U}}_b$ is contractible.*

Proof. As in the previous proof, we write $C = \{c_0, c_1, \dots, c_m\}$ with

$$c_0 \prec c_1 \prec c_2 \prec \dots \prec c_m \preccurlyeq c_0\delta$$

and assume without loss of generality that $c_0 = 1$.

The case $m = 0$ is Lemma 10.5.

Assume that $m \geq 1$. Let $T = (y, z, L)$. The point represented by T lies in $\bigcap_{b \in C} \widehat{\mathcal{U}}_b$ if and only if there exists $L_1, \dots, L_m$ with $0 < L_1 < \dots < L_m < L$ such that, for all i , $T_i := (y, z, L_i)$ is a tunnel representing c_i . Given $y \in Y$, it is possible to find such $L_1, \dots, L_m$ if and only if $y \in Y_C$. This justifies:

$$\bigcap_{b \in C} \widehat{\mathcal{U}}_{b,y} \neq \emptyset \Leftrightarrow y \in Y_C.$$

For a given $y \in Y_C$, let us study the intersection $\bigcap_{b \in C} \widehat{\mathcal{U}}_{b,y}$. Let $(x_1, \dots, x_k)$ be the ordered support of $LL(y)$. Let $A_0, \dots, A_m$ be the associated partition of $LL(y)$, defined in Proposition 9.5. Let

$$\begin{aligned} R_+(y, C) &:= \max\{\text{re}(z) \mid z \in A_m\}, \\ R_-(y, C) &:= \min\{\text{re}(z) \mid z \in A_1\}, \end{aligned}$$


$$I_+(y, C) := \min\{\operatorname{im}(z) \mid z \in A_1 \cup \dots \cup A_m\},$$

and

$$I_-(y, C) = \sup\{\operatorname{im}(z) \mid z \in A_0 \text{ and } R_- \leq \operatorname{re}(z) \leq R_+\}.$$

We have $I_-(y, C) < I_+$. It may happen that $I_-(y, C) = -\infty$.

We illustrate this on a picture. In the example, the support of $LL(y)$ is $x_1, \dots, x_6$ and $C = \{1 \prec s_2 \prec s_2 s_4\}$, where $(s_1, \dots, s_6) = \operatorname{lbl}(y)$. The support of A_1 is x_2 and the support of A_2 is x_4 . The remaining points are in A_0 . The lines $\operatorname{im}(z) = I_-(y, C)$ and $\operatorname{im}(z) = I_+(y, C)$ are represented by full lines. A semitunnel (y, z, L) representing a point in $\bigcap_{b \in C} \hat{\mathcal{U}}_{b,y}$ must cross the intervals I_2 and I_4 represented by dotted lines. One must have $\operatorname{re}(z) < R_-(y, C)$ and $I_-(y, C) < \operatorname{im}(z) < I_+(y, C)$; the final point $z + L$ must satisfy $\operatorname{re}(z + L) > R_+(y, C)$. This final point may be any complex number z' in the rectangle $\operatorname{re}(z') > R_+(y, C)$ and $I_-(y, C) < \operatorname{im}(z') < I_+(y, C)$, except the points on the closed horizontal half-line to the right of x_6 (indicated by a dashed line), which cannot be reached.



Generalising the example, one shows that $\bigcap_{b \in C} \hat{\mathcal{U}}_b$ may be identified with

$$\bigcup_{y \in Y_C} E(y, C),$$

where $E(y, C)$ is the open rectangle of $\mathbb{C}$ defined by

$$\operatorname{re}(z) > R_+(y, C), \quad I_-(y, C) < \operatorname{im}(z) < I_+(y, C),$$

from which have been removed the possible points of A_0 and the horizontal half-lines to their rights.

Let $\overline{E}(y, C)$ be the rectangle

$$\operatorname{re}(z) \geq R_+(y, C), \quad I_-(y, C) < \operatorname{im}(z) < I_+(y, C)$$

from which have been removed the possible points of A_0 and the horizontal half-lines to their rights.

A homotopy argument similar to the one used in the proof of Lemma 10.5 (or possibly a nicer argument from semialgebraic geometry) shows that $\bigcup_{y \in Y_C} E(y, C)$ and $\bigcup_{y \in Y_C} \overline{E}(y, C)$ are homotopy equivalent. The latter may be retracted to the union of open intervals $\bigcup_{y \in Y_C} (I_-(y, C), I_+(y, C))$ (on each rectangle, the retraction is

$$\overline{E}(y, C) \times [0, 1] \rightarrow \overline{E}(y, C), (z, t) \mapsto R_+(y, C) + t(\operatorname{re}(z) - R_+(y, C)) + \sqrt{-1} \operatorname{im}(z)).$$

The union $\bigcup_{y \in Y_C} (I_-(y, C), I_+(y, C))$ may be retracted to $\bigcup_{y \in Y_C} [I_0(y, C), I_+(y, C))$, where $I_0(y, C) := \max(\frac{I_-(y, C) + I_+(y, C)}{2}, I_+(y, C) - 1)$. The latter space is a fibre bundle over Y_C , with fibers intervals. Since the base space Y_C is contractible (Proposition 9.3), this fibre bundle is contractible. So is $\bigcap_{b \in C} \hat{\mathcal{U}}_b$. $\square$

11. CENTRALISERS OF REGULAR ELEMENTS

This section is devoted to the proof of Theorem 0.3. It relies on algebraic tools explained in [5], that provide us with a categorical version of the dual braid monoid with sufficient symmetries. Once one realises that such an object exists, it becomes pretty clear that proving Theorem 0.3 essentially amounts to adapting the geometric notions from the previous sections to this categorical setting. This is precisely what is done here, and it is recommended to read [5] first to get a better understanding of the conceptual framework.

Proving Theorem 0.3 involves going again through all the steps of our proof of Theorem 0.2, starting from the notion of *tunnel*, and finding suitable replacements adapted to the relative situation of Theorem 1.9. Most of the topological arguments are similar to that of the previous five sections, and actually easier.

Consider the $\mathbb{C}^\times$ -action by scalar multiplication on V . Since $W \setminus V$ and Y are defined by graded subalgebras of $\mathcal{O}_V$, we obtain quotient actions of $\mathbb{C}^\times$ on these varieties. There is also a natural $\mathbb{C}^\times$ -action on the configuration space E_n , defined by

$$\lambda \cdot \{x_1, \dots, x_n\} := \{\lambda x_1, \dots, \lambda x_n\}.$$

Note that we have to be more careful than in the previous sections, and remember that E_n consists of *centered* configurations, since the constructions involved here are not invariant by translation.

Let $d \in \mathbb{Z}_{\geq 1}$. Our goal is to understand the homotopy type of

$$(W \setminus V^{\text{reg}})^{\mu_d}.$$

We want to prove that, if it is not empty, then it is a $K(\pi, 1)$. In view of Theorem 1.9, this amounts to proving Theorem 0.3.

If d divides $h = d_n$, then $(W \setminus V^{\text{reg}})^{\mu_d}$ is again the regular orbit space of a well-generated complex reflection group (because of Theorem 1.9 (2), the centraliser W' is again a duality group, thus is well-generated). Because we have already proved the $K(\pi, 1)$ property in this case, it is enough to work under the following assumption:

Until the end of this section, we assume that d does not divide h .

We also set

$$h' := \frac{h}{h \wedge d} \quad \text{and} \quad d' := \frac{d}{h \wedge d}.$$

11.1. μ_d -action and Lyashko-Looijenga coverings.

Lemma 11.1. *For all $y \in Y$, for all $\lambda \in \mathbb{C}^\times$, we have $LL(\lambda y) = \lambda^h LL(y)$.*

Proof. Easy. □

Definition 11.2. We denote by $E_{n,\circ}$ the subspace of E_n consisting of centered configurations $\{x_1, \dots, x_n\}$ with $|x_1| = \dots = |x_n| = 1$. We set $Y_\circ := LL^{-1}(E_{n,\circ})$.

These spaces are stable under μ_d -action.

We set $V_\circ$ to be the intersection of the preimage of $Y_\circ$ with the subvariety defined by $f_n = 0$. It is clear that the natural map

$$W \setminus V_\circ \rightarrow Y_\circ$$

is a μ_d -equivariant homeomorphism.

Convention 11.3. We use this homeomorphism to identify $W \setminus V_\circ$ with $Y_\circ$. In particular, certain elements of B may be represented by paths in $Y_\circ$.

Lemma 11.4. *The space $(W \setminus V_\circ^{\text{reg}})^{\mu_d}$ is a deformation retract of $(W \setminus V^{\text{reg}})^{\mu_d}$. In particular, $(W \setminus V^{\text{reg}})^{\mu_d}$ is homotopy equivalent to $Y_\circ^{\mu_d}$.*

Proof. Let $E_{n,\times}$ be the subvariety of E_n consisting of centered configurations $\{x_1, \dots, x_n\}$ with $x_1, \dots, x_n \in \mathbb{C}^\times$ and set $Y_\times := LL^{-1}(E_{n,\times})$.

Consider a point in $W \setminus V$, represented by its coordinates

$$(x, y) \in \mathbb{C} \times Y.$$

If the point lies in $(W \setminus V)^{\mu_d}$, we must have $e^{2\pi\sqrt{-1}\frac{h}{d}}x = x$ and, since we have assumed that d does not divide h , $x = 0$. If the point lies in $(W \setminus V^{\text{reg}})^{\mu_d}$, this implies that $0 \notin LL(y)$. One concludes that $(W \setminus V^{\text{reg}})^{\mu_d}$ is isomorphic to $Y_\times$. Using the standard retraction of $\mathbb{C}^\times$ onto the unit circle and Lemma 9.4, one easily constructs a retraction of $Y_\times$ onto $Y_\circ$. $\square$

11.2. Circular tunnels. Contrary to LL , the label map lbl does not behave nicely with respect to the μ_d -action. Let us replace it with a more suitable notion. First, we need to find a good basepoint, and to replace tunnels by *circular tunnels*:

Definition 11.5. We set $\mathcal{V} := \{y \in Y_\circ \mid 0 \in U_y\}$.

In other words, y lies in $\mathcal{V}$ if and only if $LL(y)$ does not contain any point $z \in \mathbb{C}$ with $\text{re}(z) = 0$ and $\text{im}(z) > 0$.

Via the identification of $W \setminus V_\circ$ with $Y_\circ$, $\mathcal{V}$ corresponds to $\mathcal{U} \cap (W \setminus V_\circ^{\text{reg}})$. This subspace plays for $Y_\circ$ the same role as $\mathcal{U}$ was playing for $W \setminus V^{\text{reg}}$. One easily sees that it is contractible.

Definition 11.6. Let $y \in \mathcal{V}$ and $L \in [0, 2\pi]$. Let $y' := e^{\sqrt{-1}\frac{L}{h}}$. The *circular semitunnel of length L from y to y'* is the path

$$\begin{aligned} \gamma_{y,L} : [0, 1] &\rightarrow Y_\circ \\ t &\mapsto e^{\sqrt{-1}\frac{tL}{h}} y \end{aligned}$$

If $y' \in \mathcal{V}$, we say that $\gamma_{y,L}$ is a *circular tunnel*.

When $\gamma_{y,L}$ is a circular tunnel, it represents an element

$$b_{y,L} \in B.$$

One easily sees that $b_{y,L} \in M$. The condition $L \in [0, 2\pi]$ was chosen to ensure that $b_{y,L}$ is actually a simple element, as we will observe shortly.

Lemma 11.7. *For all $y \in \mathcal{V}$, $b_{y,2\pi} = \delta$.*

Proof. This is a consequence of the discussion following Definition 6.9. $\square$


Let $y \in \mathcal{V}$. Let $\{x_1, \dots, x_k\}$ be the support of $LL(y)$. Since it is included in the unit circle, we have

$$\{x_1, \dots, x_k\} = \{e^{\sqrt{-1}(\frac{\pi}{2}-\theta_1)}, e^{\sqrt{-1}(\frac{\pi}{2}-\theta_2)}, \dots, e^{\sqrt{-1}(\frac{\pi}{2}-\theta_k)}\}$$

for some sequence $(\theta_1, \theta_2, \dots, \theta_k)$ of real numbers, which is uniquely defined if require (which we naturally do) that

$$0 < \theta_1 < \theta_2 < \dots < \theta_k < 2\pi.$$

We call say that $(\theta_1, \theta_2, \dots, \theta_k)$ is the *cyclic support* of $LL(y)$. We have the corresponding sequence of *cyclic multiplicities* $(n_1, \dots, n_k)$.



By convention, set $\theta_0 := 0$. Let ε be a small enough positive real number. For $i = 1, \dots, k$, let γ_i be the circular semitunnel of length $\theta_i - \theta_{i-1}$ from $e^{\sqrt{-1}\frac{\theta_{i-1}+\varepsilon}{h}}y$ to $e^{\sqrt{-1}\frac{\theta_i+\varepsilon}{h}}y$. It is clear that, if ε is small enough, γ_i is actually a circular tunnel, and that the element

$$c_i \in M$$

represented by γ_i does not depend on the choice of ε .

Definition 11.8. The *cyclic label* of y is the sequence


$$cbl(y) := (c_1, \dots, c_k)$$

as constructed above.

By construction, one has

$$c_1 \dots c_k = \delta.$$

It is easily seen that the cyclic label consists of simple elements: up to replacing y by some $e^{\sqrt{-1}\frac{\theta_i+\varepsilon}{h}}y$, it is enough to show that c_1 is simple. In the above example, a tunnel representing c_1 is as follows:



The following definition is an analog of Definition 7.8:

Definition 11.9. Let $x \in E_n$ be such that the support of x is in the unit circle. Let $(\theta_1, \dots, \theta_k)$ and $(n_1, \dots, n_k)$ be the cyclic support and multiplicities of x . Let $\underline{c}$ be a finite sequence of simple elements. We say that x and $\underline{c}$ are *cyclically compatible* if the following conditions are satisfied:

- $\underline{c}$ consists of k terms $c_1, \dots, c_k$,
- for all i , $l(c_i) = n_i$,
- $c_1 \dots c_k = \delta$.

We have the following cyclic analog of Theorem 7.9:

Proposition 11.10. *The map $LL \times cbl : Y_\circ \rightarrow E_{n,\circ} \times S^*$ is injective, and its image is the subset of $E_{n,\circ} \times S^*$ consisting of cyclically compatible pairs $(x, \underline{c})$.*

Sketch of proof. The result is an easy consequence of Theorem 7.9, once it has been observed that, for a given y with known $LL(y)$, the value of $cbl(y)$ may be recovered from that of $lbl(y)$, and vice-versa. We leave to the reader to describe the correspondence between $cbl(y)$ and $lbl(y)$, and to check that it transforms the compatibility conditions of Definition 7.8 into that of 11.9. $\square$

Let us now explain how $clbl(y)$ varies along a circular tunnel. Let $y \in \mathcal{V}$. Let $(\theta_1, \dots, \theta_k)$ be the cyclic support of $LL(y)$.

Let $\theta \in [0, 2\pi]$. If θ is one the θ_i , then the circular semitunnel of length θ from y to $y' := e^{\sqrt{-1}\frac{\theta}{h}}y$ is not a tunnel. Otherwise, it is a tunnel and the cyclic label of y' may be computed from $(c_1, \dots, c_k) := clbl(y)$ as follows:

If $0 \leq \theta < \theta_1$, the tunnel is within $\mathcal{V}$ and $clbl(y') = clbl(y)$.

If $\theta_1 < \theta < \theta_2$, then the tunnel crosses exactly once the complement of $\mathcal{V}$ and we have

$$clbl(y') = (c_2, \dots, c_k, c_1^\delta),$$

where $c_1^\delta := \delta^{-1}c_1\delta$. Indeed, the first $k-1$ terms of $clbl(y')$ are, by construction, the last $k-1$ terms of $clbl(y)$; the last term is then uniquely determined by the condition that the product should be δ .

More generally, if $\theta_i < \theta < \theta_{i+1}$, we have

$$clbl(y') = (c_{i+1}, \dots, c_n, c_1^\delta, \dots, c_i^\delta).$$

11.3. m -cyclic labels. Although the map $LL \times clbl$ is already more effective than $LL \times lbl$ at handling $Y_\circ^{\mu_d}$, it is not yet entirely practical, since we want to avoid discussing according to the particular values of $\theta_1, \dots, \theta_k$, which of course depend on y . It is convenient to introduce of the following variant of $clbl$:

Definition 11.11. Let $y \in Y_\circ$ and let m be a positive integer. Assume that

$$\{y, e^{\frac{2i\pi}{h}\frac{1}{m}}y, e^{\frac{2i\pi}{h}\frac{2}{m}}y, \dots, e^{\frac{2i\pi}{h}\frac{m-1}{m}}y\} \subseteq \mathcal{V}.$$

The m -cyclic label of y is the sequence

$$clbl_m(y) := (f_1, \dots, f_m)$$

where $f_i \in S$ is represented by the circular tunnel of length $\frac{1}{m}$ from $e^{\frac{2i\pi}{h}\frac{i-1}{m}}y$ to $e^{\frac{2i\pi}{h}\frac{i}{m}}y$.

When it is defined, the m -cyclic label may be compared with

$$clbl(y) = (c_1, \dots, c_k).$$

Let $(\theta_1, \dots, \theta_k)$ be the cyclic support. Let

$$0 = j_0 \leq j_1 \leq \dots \leq j_{m-1} \leq j_m = k$$


be such that, for all $j \in \{1, \dots, k\}$,

$$\theta_j \in [\frac{j-1}{m}, \frac{j}{m}] \Leftrightarrow j_{i-1} < j \leq j_i.$$

Then

$$clbl_m(y) = (c_1 \dots c_{j_1}, c_{j_1+1} \dots c_{j_2}, \dots, c_{j_{m-1}+1} \dots c_k).$$

Let us illustrate this with $m = 4$:



The function $clbl_m$ is coarser than $clbl$, and is not able to distinguish whether the intersection of $LL(y)$ with a given cone of angle $2\pi/m$ consists of distinct points. However, if we know $clbl_m(\zeta y)$ for all ζ in the unit circle, it is possible to recover $clbl(y)$.

11.4. A relative dual categorical Garside structure for W' . Denote by D_m the set of factorisations of δ into m simple terms, some of which may be trivial. By the above discussion,

$$clbl_m(y) \in D_m.$$

As explained in [5], one may define a m -divided Garside category of (M, δ) , denoted by M_m , whose object set is D_m .

One may think of M_m as a variant of M that retains its most important features (including the possibility to construct a nice simplicial $K(B, 1)$) but whose automorphism group is artificially enlarged (at the expense of increasing the number of objects). The case $m = 1$ corresponds to M itself: a monoid is nothing but a small category with a single object, the elements of the monoid being interpreted as the endomorphisms of the object. So far, we have only mentioned the objects of M_m . Morphisms will later be understood in terms of circular tunnels.

The category M_m admits an automorphism denoted by ϕ_m which acts on objects by

$$(f_1, \dots, f_m) \cdot \phi_m := (f_2, \dots, f_m, f_1^\delta)$$

(we like to view it as acting *on the right*, since it is a sort of “right conjugacy”).

Lemma 11.12. *Let m be a positive integer. Let $y \in \mathcal{V}_m$. Then*

$$clbl_m(\zeta_{mh}y) = clbl_m(y) \cdot \phi_m.$$

Proof. It follows from our above discussions about $clbl(e^{\sqrt{-1}\frac{\theta}{h}}y)$ and about the relationship between $clbl$ and $clbl_m$. $\square$

Lemma 11.13. *If $y \in \mathcal{V}^{\mu_d}$, then $clbl_{d'}(y) \in M_{d'}^{\phi_{d'}^{h'}}$. The following assertions are equivalent:*

- (i) *The integer d is regular.*
- (ii) *The category $M_{d'}$ admits $\phi_{d'}^{h'}$ -invariant objects.*
- (iii) *The category M_d admits ϕ_d^h -invariant objects.*

Proof. If $y \in \mathcal{V}^{\mu_d}$, then $clbl_{d'}(y) = clbl_{d'}(\zeta_d y) = clbl_{d'}(\zeta_{dh'}^{h'} y) = clbl_{d'}(\zeta_{d'h}^{h'} y)$. Using Lemma 11.12, this is equal to $clbl_{d'}(y) \cdot \phi_{d'}^{h'}$. We have proved the first statement, which itself implies (i) $\Rightarrow$ (ii). Conversely, given any $\phi_{d'}^{h'}$ -fixed object $(c_1, \dots, c_{d'})$ of $M_{d'}$, we may use Proposition 11.10 to find $y \in Y_o^{\mu_d}$ with $clbl_{d'}(y) = (c_1, \dots, c_{d'})$.

The equivalence (i) $\Leftrightarrow$ (iii) can be proved similarly (an alternate way to conclude is via the equivalence (ii) $\Leftrightarrow$ (iii), which is a general property of divided Garside categories, see [5, Proposition 9.8]). $\square$

Definition 11.14. The *reduced dual Garside category* for W' relative to the pair (W, W') is the fixed subcategory

$$M' := M_{d'}^{\phi_{d'}^{h'}}.$$

We denote by G' the groupoid obtained by formally inverting the morphisms in M' . We denote by O' the common object set to M' and G' (the elements of O' are $\phi_{d'}^{h'}$ -invariant factorisations of δ into d' simple elements of M).

The centraliser W' (see Theorem 1.9) may be badly-generated, but the category M' provides a substitute for the dual braid monoid.

The following theorem is an analog for W' of the “Garside part” of Theorem 8.2:

Theorem 11.15. *The category M' is a Garside category. As a consequence, G' is a Garside groupoid.*

Proof. See [5]: because M is a Garside monoid, $M_{d'}$ is a Garside category, and the fixed subcategory by an automorphism of a Garside category is a Garside category. $\square$

Both M' and G' can be viewed as subcategories of the fundamental groupoid of $Y_o^{\mu_d}$, as it will appear later.

Proposition 11.16. *The map*

$$cbl_{d'} : \mathcal{V}^{\mu_d} \rightarrow O'$$

is surjective and its fibers are the connected components of $\mathcal{V}^{\mu_d}$. These connected components are contractible.

Proof. Note first that if $y \in \mathcal{V}^{\mu_d}$, then $LL(y)$ contains no point with argument in $\frac{\pi}{2} + \frac{2\pi}{d'}\mathbb{Z}$. Indeed, by Lemma 11.1, $LL(y)$ is ζ_d^h -invariant, thus it is $\zeta_{d'}$ -invariant; the condition $y \in \mathcal{V}$ forces that $LL(y)$ contains no point with argument $\frac{\pi}{2}$; the conclusion follows.

In particular, the value of $cbl_{d'}$ cannot change along a continuous path in $\mathcal{V}^{\mu_d}$. Each connected component of $\mathcal{V}^{\mu_d}$ must be included in a fiber of $cbl_{d'}$.

Choose a real number θ in the open interval $]\frac{\pi}{2} - \frac{2\pi}{d'}, \frac{\pi}{2}[$. Consider the multiset

$$x_0 := \left\{ \underbrace{e^{\sqrt{-1}\theta}, \dots, e^{\sqrt{-1}\theta}}_{\frac{n}{d'} \text{ copies}}, \underbrace{e^{\sqrt{-1}(\theta + \frac{2\pi}{d'})}, \dots, e^{\sqrt{-1}(\theta + \frac{2\pi}{d'})}}_{\frac{n}{d'} \text{ copies}}, \dots, \underbrace{e^{\sqrt{-1}(\theta + \frac{2\pi(d'-1)}{d'})}, \dots, e^{\sqrt{-1}(\theta + \frac{2\pi(d'-1)}{d'})}}_{\frac{n}{d'} \text{ copies}} \right\}.$$

One can clearly deformation retract $\mathcal{V}^{\mu_d}$ onto $\mathcal{V}^{\mu_d} \cap LL^{-1}(x_0)$.

Claim: the map $cbl_{d'} : \mathcal{V}^{\mu_d} \cap LL^{-1}(x_0) \rightarrow O'$ is a bijection. Indeed, since x_0 contains exactly one point (without multiplicity) in each cone of angle $\frac{2\pi}{d'}$, the maps $cbl_{d'}$ and cbl coincide on $\mathcal{V}^{\mu_d} \cap LL^{-1}(x_0)$. Using Proposition 11.10, we conclude that $cbl_{d'}$ induces a bijection from $\mathcal{V} \cap LL^{-1}(x_0)$ to the set of factorisations of δ in d' simple factors of individual length $\frac{n}{d'}$. By Lemma 11.12, it is obvious that $y \in \mathcal{V} \cap LL^{-1}(x_0)$ is μ_d -invariant if and only if cbl_m is an object of M' .

Combining the claim and the fact that $\mathcal{V}^{\mu_d}$ deformation retracts onto $\mathcal{V}^{\mu_d} \cap LL^{-1}(x_0)$, we see that the map $cbl_{d'}$ induces a bijection between connected components of $\mathcal{V}^{\mu_d}$ and objects of M' . $\square$

11.5. An interpretation of the groupoid G' . Consider the category

$$B' := \pi_1(Y_o^{\mu_d}, \pi_0(\mathcal{V}^{\mu_d})),$$

whose objects are connected components of $\mathcal{V}^{\mu_d}$ and morphisms are relative homotopy classes of paths in Y_o with endpoints in $\mathcal{V}^{\mu_d}$. The category structure is provided by the fact that the connected components are simply connected: up to homotopy, there is a natural way to concatenate a path arriving somewhere in a connected component to a path leaving from somewhere else in the same component (this is the groupoid version of the trick that consists of replacing a basepoint by a simply-connected subspace). Equivalently, although it would have been less convenient, we could have worked with one favorite actual basepoint chosen in each connected component – the advantage of the latter viewpoint is that it identifies B' with a full subcategory of the fundamental groupoid of $Y_o^{\mu_d}$.

The following lemma simply says that B' is a “multi-basepoint” version of $B(W')$:

Lemma 11.17. *The groupoid B' is equivalent as a category to $B(W')$, the braid group of W' (defined as the fundamental group of $(W \setminus V^{\text{reg}})^{\mu_d}$).*

Proof. By Lemma 11.4, $Y_o^{\mu_d}$ is homotopy equivalent to the regular orbit space $(W \setminus V^{\text{reg}})^{\mu_d}$, thus their fundamental groupoids are equivalent. Since both spaces are path connected, their

fundamental groupoids are connected and equivalent to any of their non-empty full subcategories. $\square$

We will prove that B' and G' are isomorphic.

Using Proposition 11.16, we may already observe that $clbl_{d'}$ induces a bijection from the object set of B' to the object set O' of G' . To simplify notations, we identify the object set of B' with O' .

Definition 11.18. An element of $s \in B'$ is said to be *simple* if there exists $y \in \mathcal{V}^{\mu_d}$ and $L \in [0, \frac{2\pi}{d'}]$ such that $e^{\sqrt{-1}\frac{L}{h}}y \in \mathcal{V}^{\mu_d}$, and s is represented by the circular tunnel of length L starting at y .

We denote by S' the set of simple elements in B' . It is part of an oriented graph structure

$$S' \begin{matrix} \xrightarrow{\partial_0} \\ \xRightarrow{\partial_1} \end{matrix} O'$$

where ∂_0 is the source map and ∂_1 is the target map.

The category M' comes equipped its own notion of *simple elements*, which is part of its natural *categorical Garside structure*. The simple elements form a graph $\Sigma' \begin{matrix} \xrightarrow{\partial_0} \\ \xRightarrow{\partial_1} \end{matrix} O'$ with the following properties (see [5]):

- It is a subgraph of (the oriented graph underlying to) the category M' (in particular, ∂_0 is the source map and ∂_1 is the target map).
- Edges are $2d'$ -tuples

$$F = (f'_1, f''_1, f'_2, f''_2, \dots, f'_{d'}, f''_{d'})$$

of elements in S such that $f'_1 f''_1 f'_2 f''_2 \dots f'_{d'} f''_{d'} = \delta$. The source and target of such an F are as follows:

$$\partial_0(F) = (f'_1 f''_1, f'_2 f''_2, \dots, f'_{d'} f''_{d'})$$

$$\partial_1(F) = (f''_1 f'_2, f''_2 f'_3, \dots, f''_{d'} f'_1{}^\delta).$$

- The category M' is generated by Σ' .
- Let $(f_1, \dots, f_{d'})$ be an object. Choose factorisations $f_i = f'_i f''_i f'''_i$ of each f_i into three simples of M . We obtain three edges $F, G, H \in \Sigma'$ as follows:

$$\begin{array}{ccc} (f'_1 f''_1 f'''_1, f'_2 f''_2 f'''_2, \dots, f'_{d'} f''_{d'} f'''_{d'}) & & \\ & \searrow F = (f'_1, f''_1 f'''_1, f'_2, f''_2 f'''_2, \dots, f'_{d'}, f''_{d'} f'''_{d'}) & \\ H = (f'_1 f''_1, f'_1 f'''_1, f'_2 f''_2, f'_2 f'''_2, \dots, f'_{d'} f''_{d'}, f'_{d'} f'''_{d'}) & & (f''_1 f'''_1 f'_2, f''_2 f'''_2 f'_3, \dots, f''_{d'} f'''_{d'} f'_1{}^\delta) \\ & \swarrow G = (f''_1, f'''_1 f'_2, f''_2, f'''_2 f'_3, \dots, f''_{d'}, f'''_{d'} f'_1{}^\delta) & \\ & (f'''_1 f'_2 f''_2, f'''_2 f'_3 f''_3, \dots, f'''_{d'} f'_1{}^\delta f''_1{}^\delta) & \end{array}$$

Then the relation $FG = H$ holds in M' . Moreover, these are defining relations for M' : the category is defined as the quotient of the free category of walks on Σ' by the smallest congruence $\sim$ such that $FG \sim H$ for all triples (F, G, H) obtained as above.

Theorem 11.19. *There exists a unique bijection*

$$\Theta : \Sigma' \xrightarrow{\sim} S'$$

such that $\partial_0 \circ \Theta = \partial_0$ and $\partial_1 \circ \Theta = \partial_1$. This bijection induces category isomorphisms

$$\Theta : M' \xrightarrow{\sim} \langle S' \rangle_+$$

(where $\langle S' \rangle_+$ is the subcategory of B' generated by S') and

$$\Theta : G' \xrightarrow{\sim} B'.$$

In light of Lemma 11.17, the theorem should be understood as an analog for the centraliser W' of Theorem 8.2.

Proof. Let $s \in S'$, let y and L be as in Definition 11.18. Let γ be the circular tunnel of length L starting at y . It represents s .

Let

$$(c_1, \dots, c_k) := clbl(y),$$

$$(f_1, \dots, f_{d'}) := clbl_{d'}(y).$$

By μ_d -invariance, it is clear that k is a multiple of d' , say $k = ed'$, and that, for all i ,

$$f_i = c_{(i-1)e+1} \cdots c_{ie}$$

When t varies from 0 to 1, some points in $LL(\gamma(t))$ may cross the vertical half-line of argument $\frac{\pi}{2}$. These points were originally labelled by $(c_1, \dots, c_j)$, where j is an integer in $[0, e]$ (the integer j increases with L ; if $L = 0$, then $j = 0$; if $L = \frac{2\pi}{d'}$, then $j = e$).

For all j , set

$$f'_i := c_{(i-1)e+1} \cdots c_{(i-1)e+j}$$

and

$$f''_i := c_{(i-1)e+j+1} \cdots c_{ie}.$$

In other words, in our previous expression of f_i as a product of e terms in $clbl(y)$, we take f'_i to be the product of the first j terms, and f''_i to be the product of the remaining terms. In particular,


$$f_i = f'_i f''_i.$$

A direct computation shows that

$$clbl_{d'}(e^{\sqrt{-1}\frac{L}{h}} y) = clbl_{d'}(\gamma(1)) = (f'_1 f'_2, f''_2 f'_3, \dots, f''_{d'} f'_1).$$

Moreover, the factorisations $f_i = f'_i f''_i$ are uniquely determined by s . This defines a map $\Theta' : S' \rightarrow \Sigma'$, compatible with ∂_0 and ∂_1 .

When $e \leq 2$, we say that y is *elementary*. The case $e = 2$ looks as follows:



The case $e = 1$ happens when either $f'_1 = \cdots = f'_{d'} = 1$ (thus s is the identity morphism of its source) or $f''_1 = \cdots = f''_{d'} = 1$ (s is the *Garside element* starting at its source).

In the general case when y is not assumed to be elementary, we may always replace it by another $\tilde{y} \in \mathcal{V}^{\mu_d}$ such that $\tilde{y}$ is elementary and s is represented by the circular tunnel of length L starting at $\tilde{y}$. This is done by collapsing the different points in the support of y corresponding

to $c_{(i-1)e+1}, \dots, c_{(i-1)e+j}$ (resp. $c_{(i-1)e+j+1}, \dots, c_{ie}$) into a single point (or no point when $j = 0$, resp. when $j = e$) and applying an obvious “circular Hurwitz rule”.

We now construct the map $\Sigma' \rightarrow S'$. Given any object $(f_1, \dots, f_{d'}) \in O'$ and any factorisation $(f'_1, f''_1, \dots, f'_{d'}, f''_{d'}) \in \Sigma'$ with source $(f_1, \dots, f_{d'})$, we may apply Proposition 11.10 to find $y \in \mathcal{V}^{\mu_d}$ and $L \in [0, \frac{2\pi}{d}]$ such that:

- $clbl_{d'}(y) = (f_1, \dots, f_{d'})$,
- y is elementary,
- $e^{\sqrt{-1}\frac{L}{h}}y \in \mathcal{V}^{\mu_d}$ and
- $clbl_{d'}(e^{\sqrt{-1}\frac{L}{h}}y) = (f'_1 f'_2, f''_2 f'_3, \dots, f''_{d'} f'_1)$.

Moreover, one easily sees that the simple element s associated to y and L is independent of the choice of specific y and L satisfying the above conditions (to see this, use Proposition 11.10 to describe all possible choices and observe that they only differ by small angles that have no impact on s).

This defines a map $\Theta : \Sigma' \rightarrow S'$ compatible with ∂_0 and ∂_1 . By construction, it is clear that $\Theta' \circ \Theta = 1_{\Sigma'}$. It is also clear that, if $s \in S'$ is represented by a circular tunnel starting at an elementary $y \in \mathcal{V}^{\mu_d}$, $\Theta \circ \Theta'(s) = s$. As we have noted, any $s \in S'$ may be represented by a circular tunnel starting at an elementary point. Thus Θ and Θ' are inverse bijections.

To prove that Θ is the *unique* $\Sigma' \xrightarrow{\sim} S'$ compatible with ∂_0 and ∂_1 , it suffices to show that, for any $(f_1, \dots, f_{d'}), (g_1, \dots, g_{d'}) \in O'$, there is *at most* one arrow $F = (f'_1, f''_1, f'_2, f''_2, \dots, f'_{d'}, f''_{d'})$ in Σ' with source $(f_1, \dots, f_{d'})$ and target $(g_1, \dots, g_{d'})$. Clearly, it suffices to show that f'_1 is uniquely determined by $(f_1, \dots, f_{d'})$ and $(g_1, \dots, g_{d'})$. The natural injection

$$Y_o^{\mu_d} \simeq (W \setminus V_o^{\text{reg}})^{\mu_d} \hookrightarrow (W \setminus V^{\text{reg}})^{\mu_d} \hookrightarrow W \setminus V^{\text{reg}}$$

maps $\mathcal{V}^{\mu_d}$ into the “basepoint” $\mathcal{U}$, thus we have a natural functor $\psi : B' \rightarrow B$. A direct computation using Hurwitz rule shows that $\psi(\Theta(F)) = f'_1$ (the argument is the same that we used to show that the cyclic label consists of elements of S ; see the illustration above Definition 11.9). This proves unicity.

Let us now prove that Θ extends to a morphism $M' \rightarrow \langle S' \rangle_+$, *i.e.*, that whenever a defining relation $FG = H$ holds in Σ' , we have $\Theta(F)\Theta(G) = \Theta(H)$. The relation corresponds to a factorisation

$$R := (f'_1, f''_1, f'''_1, f'_2, f''_2, f'''_2, \dots, f'_{d'}, f''_{d'}, f'''_{d'})$$

of δ as a product of $3d'$ elements of S . Applying Proposition 11.10, we may find $y \in \mathcal{V}^{\mu_d}$ such that $clbl(y) = R$. For suitable $L, L' \in [0, \frac{2\pi}{d}]$, $\Theta(F)$ (resp. $\Theta(G)$, resp. $\Theta(H)$) is represented by the circular tunnel of length L (resp. L' , resp. $L + L'$) starting at y (resp. $e^{\sqrt{-1}\frac{L}{h}}y$, resp. y). The conclusion follows.

Because the groupoid G' is the localisation of M' , the functor $\Theta : M' \rightarrow \langle S' \rangle_+$ uniquely extends to a functor $\Theta : G' \rightarrow B'$. To prove that the latter is an isomorphism, we have to show that all elements of B' are products of simples and inverses of simples, and that the relations $\Theta(F)\Theta(G) = \Theta(H)$ are defining relations for B' . We follow Brieskorn’s strategy from [15] and stratify $Y_o^{\mu_d}$ according to the multiplicity of $e^{\sqrt{-1}\frac{\pi}{2}}$ in the support of $LL(y)$:

$$Y_o^{\mu_d} = X_0 \sqcup X_1 \sqcup \dots \sqcup X_{n/d'}$$

where $X_k := \{y \in Y_o^{\mu_d} | e^{\sqrt{-1}\frac{\pi}{2}}$ has multiplicity k in $LL(y)\}$ (because of the μ_d -invariance, the multiplicity of $LL(y)$ is constant on $e^{\sqrt{-1}\frac{\pi}{2}}\mu_{d'}$, thus cannot be larger than n/d'). The open stratum X_0 is precisely our “multibasepoint” $\mathcal{V}^{\mu_d}$. Each stratum X_k has real codimension k in $Y_o^{\mu_d}$. A standard general position argument (as in [15]) shows that the natural functor

$$\pi_1(X_0 \sqcup X_1, X_0) \rightarrow B'$$

is surjective (“full”) and that the natural functor

$$\pi_1(X_0 \sqcup X_1 \sqcup X_2, X_0) \rightarrow B'$$

is an isomorphism. One deduces that B' is generated (as a groupoid) by the simples of the form $\Theta(F)$, where $F \in \Sigma'$ has a first term f'_1 of length 1. The isomorphism $\pi_1(X_0 \sqcup X_1 \sqcup X_2, X_0) \simeq B'$ shows that B' has a presentation with those generators and relations of length 2, of the form $\Theta(F)\Theta(G) = \Theta(F')\Theta(G')$. These relations are associated with components of X_2 and follow from our relations $\Theta(F)\Theta(G) = \Theta(H)$, where H has a first term of length 2.

Because the localisation functor $M' \rightarrow G'$ is faithful (this is general property of Garside categories) and $\Theta : G' \rightarrow B'$ is an isomorphism, the functor from M' to $\langle S' \rangle_+$ must also be an isomorphism. $\square$

The theorem may be used to write down an explicit presentation for B' and $B(W')$. For example, one may prove the case G_{31} of Theorem 0.6 without any computer-assisted monodromy computations (improving the argument given at the end of Section 4). Some other applications are:

Corollary 11.20. *The braid group $B(W')$ is a weak Garside group, in the sense of [5].*

Corollary 11.21. *The categories M' and G' are connected.*

Proof. This is because B' is a connected category (see Lemma 11.17). $\square$

11.6. The universal cover. Let us fix a particular object $* \in O'$ – in light of Theorem 11.19, this amounts to choosing a preferred connected component, still denoted by $*$, of $\mathcal{V}^{\mu_d}$.

Define a simplicial complex Γ' as follows:

- The vertex set is

$$\text{Hom}_{G'}(*, -) := \bigsqcup_{o \in O'} \text{Hom}_{G'}(*, o).$$

- Two vertices $g \in \text{Hom}_{G'}(*, o)$ and $g' \in \text{Hom}_{G'}(*, o')$ span an edge if and only if $g^{-1}g' \in \Sigma'$ or $g'^{-1}g \in \Sigma'$.

The associated flag complex $\text{Flag}(\Gamma')$ is the natural generalisation to our setting of the complex described in Definition 8.4.

Because G' is a Garside groupoid with graph of simples Σ' , a generalisation of Theorem 8.5 is applicable (see [5, Corollary 7.6]). It implies that the geometric realisation of $\text{Flag}(\Gamma')$ is contractible.

Let us use this to prove that $Y_o^{\mu_d}$ is a $K(\pi, 1)$. As in Section 10, we have a model of the universal cover $\widehat{Y_o^{\mu_d}}$ where points are represented by paths starting in $*$ and ending anywhere. The open covering used in Section 10 was the $B(W)$ -orbit of a basic piece $\widehat{\mathcal{U}}_1$. Here, we need a basic piece for each element in our desired index set, the vertex set $\text{Hom}_{G'}(*, -)$ of Γ' .

The following definition makes use of the identification of (B', S') with (G', Σ') (Theorem 11.19).

Definition 11.22. Let $g \in \text{Hom}_{G'}(*, -)$. We denote by $\widehat{\mathcal{W}}_g$ the open subset of $\widehat{Y_o^{\mu_d}}$ consisting of points represented by paths of the form γT , where:

- γ is any path in $Y_o^{\mu_d}$ with source in $*$ and target in $\mathcal{V}^{\mu_d}$ and representing g .
- T is a circular semitunnel of length $< \frac{2\pi}{d}$ starting at the target of γ .

Lemma 11.23. *The family $(\widehat{\mathcal{W}}_g)_{g \in \text{Hom}_{G'}(*, -)}$ is a locally finite open covering of $\widehat{Y_o^{\mu_d}}$, whose intersections are either empty or contractible. The nerve of $(\widehat{\mathcal{W}}_g)_{g \in \text{Hom}_{G'}(*, -)}$ is $\text{Flag}(\Gamma')$.*

Theorem 0.3 follows: the lemma shows that $Y_o^{\mu_d}$ is homotopy equivalent to the realisation of flag complex of $\text{Flag}(\Gamma')$. By [5, Corollary 7.6], the latter space is a $K(\pi, 1)$. By Lemma 11.4, $Y_o^{\mu_d}$ is homotopy equivalent to $(W \setminus V^{\text{reg}})^{\mu_d}$, which coincides (Theorem 1.9) with the space $W' \setminus V'^{\text{reg}}$ involved in Theorem 0.3.

Proof. The lemma contains analogs for $(\widehat{\mathcal{W}}_g)_{g \in \text{Hom}_{G'}(*, -)}$ of various results about $(\widehat{\mathcal{U}}_b)_{b \in B}$, namely Lemma 10.5 (the sets are open and contractible), Proposition 10.6 (the nerve of the covering is $\text{Flag}(\Gamma')$) and Proposition 10.7 (non-empty intersections are contractible). The strategy is to imitate the material in Sections 9 and 10. The situation is actually much easier, because we no longer have to care about the fibers of $W \setminus V^{\text{reg}} \rightarrow Y$.

First, one defines the obvious analogs $\overline{\mathcal{W}}_g \subseteq \widehat{\mathcal{W}}_g$ and $\mathcal{W}_g \subseteq \overline{\mathcal{W}}_g$ of the subsets $\overline{\mathcal{U}}_b \subseteq \widehat{\mathcal{U}}_b$ and $\mathcal{U}_b \subseteq \overline{\mathcal{U}}_b$. One shows that $(\overline{\mathcal{W}}_g)_{g \in \text{Hom}_{G'}(*, -)}$ is a partition of $Y_o^{\mu_d}$. Imitating the proof of Lemma 10.5, one sees that each $\widehat{\mathcal{W}}_g$ is open and contractible.

Let $g_0, \dots, g_k$ be distinct elements of $\text{Hom}_{G'}(*, -)$. Assume that the intersection $\widehat{\mathcal{W}}_{g_0} \cap \dots \cap \widehat{\mathcal{W}}_{g_k}$ is non-empty and let z be a point in this intersection. We may find, for each $i \in \{0, \dots, k\}$, a path γ_i with source in $*$ and target $y_i \in \mathcal{V}^{\mu_d}$ representing g_i and $L_i \in [0, \frac{2\pi}{d}]$ such that the z is represented by each of the paths $\gamma_i T_i$, where T_i is the circular semitunnel of length L_i starting y_i . Up to reordering the g_i 's, we may assume that

$$L_0 \geq L_1 \geq \dots \geq L_k$$

(the reordering *a priori* depends on z but, as it will appear shortly, it only depends $g_0, \dots, g_k$). Since $\gamma_0 T_0, \dots, \gamma_k T_k$ represent the same point in the universal cover, they have the same target, thus

$$e^{\sqrt{-1}\frac{L_0}{h}} y_0 = \dots = e^{\sqrt{-1}\frac{L_k}{h}} y_k.$$

Consider the circular tunnel $T_{0,i}$ of length $L_0 - L_i$ starting at y_0 . Its target coincides with y_i and it represents some element $s_i \in \Sigma'$. Using again that $\gamma_0 T_0$ and $\gamma_i T_i$ represent the same point in the universal cover, we see that $\gamma_0 T_{0,i}$ and γ_i represent the same point and, in particular, that

$$g_0 s_i = g_i.$$

The same argument can be applied to show that, if $i \leq j$, there exists $t_{ij} \in \Sigma'$ such that $s_i t_{ij} = s_j$. This proves that $\{g_0, \dots, g_k\}$ is a k -simplex.

Because the groupoid G' is equipped with a canonical group morphism $G' \rightarrow \mathbb{Z}$ mapping Σ' into $\mathbb{Z}_{\geq 0}$, we observe that, for any k -simplex in $\text{Flag}(\Gamma')$, there is a unique way to label its elements $g_0, \dots, g_k$ such that, for all i , $g_0^{-1} g_i \in \Sigma'$. When this is satisfied, $i \leq j \Rightarrow g_i^{-1} g_j \in \Sigma'$.

Assume now that $\{g_0, \dots, g_k\}$ is a k -simplex in Γ' . Let us prove that $\widehat{\mathcal{W}}_{g_0} \cap \dots \cap \widehat{\mathcal{W}}_{g_k}$ is non-empty and contractible. This will complete the proof of the lemma.

We choose the labelling in such a way that each $s_i := g_0^{-1} g_i$ lies in Σ' . Because G' acts transitively on Γ' (by concatenation at the source), it suffices to work under the assumption that $g_0 = 1_*$.

Our base object $* \in O'$ is a factorisation $(f_1, \dots, f_{d'})$ of δ into elements of S . Each s_i consists of factorisations $f_j = f'_{i,j} f''_{i,j}$ for all j . We focus our attention to $j = 1$ and set $h_i := f'_{i,1}$. If $i, i' \in \{1, \dots, k\}$ are such that $i < i'$, then $s_i^{-1} s_{i'} \in \Sigma'$ and $h_i \prec h_{i'}$. We have a chain in S as follows:

$$1 \prec h_1 \prec \dots \prec h_k \preceq f_1$$

Let y_0 be a point in the base component $\mathcal{V}^{\mu_d}$ associated with $*$, *i.e.*, in the fiber of $clbl_{d'}$ over $(f_1, \dots, f_{d'})$. Let

$$(c_1, \dots, c_m) := clbl(y_0)$$

and let

$$(\theta_1, \dots, \theta_m)$$

be the cyclic support of $LL(y_0)$. We have $m = d'e$ for a certain integer e , and $f_1 = c_1 \dots c_e$. We say that y_0 is *admissible* (with respect to $(s_1, \dots, s_k)$) if, for all i , there exists an integer j_i such that

$$h_i = c_1 \dots c_{j_i}.$$

Clearly, when this happens, we have

$$0 < j_1 < \dots < j_k \leq e.$$

When y_0 is admissible, we set $L_{\min}(y_0) := \theta_{j_k}$. One has $0 < L_{\min}(y_0) < \frac{2\pi}{d'}$.

Claim: A point z lies in $\widehat{\mathcal{W}}_{g_0} \cap \dots \cap \widehat{\mathcal{W}}_{g_k}$ if and only if it is represented by a circular semitunnel whose source is an admissible point y_0 and whose length lies in $(L_{\min}(y_0), \frac{2\pi}{d'})$.

The claim is an easy consequence of our earlier discussion of necessary conditions for z to be in $\widehat{\mathcal{W}}_{g_0} \cap \dots \cap \widehat{\mathcal{W}}_{g_k}$.

From the claim, one deduces that $\widehat{\mathcal{W}}_{g_0} \cap \dots \cap \widehat{\mathcal{W}}_{g_k}$ is homotopy equivalent to the space of admissible points.

The latter space may be described thanks to Proposition 11.10. It is clearly non-empty. It deformation retracts onto the space of admissible points with $e = k$ (see above notations), which is obviously contractible. (Remark: The fact that the space of admissible points is contractible plays the role of Proposition 9.3). $\square$

12. PERIODIC ELEMENTS IN $B(W)$

As before, W is an irreducible well-generated complex reflection group, $\tau \in P(W)$ is the full-twist and $\delta \in B(W)$ is the Garside element of the dual braid monoid $M(W)$. The image of δ in W is a Coxeter element c .

Definition 12.1. An element of $B(W)$ is *periodic* if it admits a central power.

The goal of this section is to prove that the center $ZB(W)$ is cyclic, and to establish a correspondence between periodic elements in $B(W)$ and regular elements in W . As with the previous section, the real substance of the arguments lies more in the algebraic tools from [5] than in the easy geometric interpretation.

Lemma 12.2. *The intersection of the subgroup $\langle c \rangle \subseteq W$ with the interval $[1, c]$ is $\{1, c\}$.*

Proof. [27, Proposition 4.2] gives an argument for the real case that, as pointed out by J. Michel, applies verbatim. We include it for the convenience of the reader. Let $2 \leq d_1, \dots, d_n = h$ be the reflection degrees. Because c is regular, its eigenvalues are $\zeta_d^{1-d_i}$ ([41, Theorem 4.2 (v)]). Assume that some power c^k lies in $[1, c]$, thus that

$$l_R(c^k) + l_R(c^{1-k}) = n.$$

By Proposition 8.7 and Lemma 7.3,

$$l_R(c^k) = n - \#\{i | (1 - d_i)k \equiv 0[h]\}.$$

Using the same formula for c^{1-k} , we deduce that

$$\#\{i | (1 - d_i)k \equiv 0[h]\} + \#\{i | (1 - d_i)(1 - k) \equiv 0[h]\} = n.$$

For a given i , $(1 - d_i)k$ and $(1 - d_i)(1 - k)$ cannot simultaneously divide h . The identity then forces that, for all i , either $(1 - d_i)k$ or $(1 - d_i)(1 - k)$ is a multiple of h . But, when $2 \leq k \leq h - 1$, neither $(1 - h)k$ nor $(1 - h)(1 - k)$ is a multiple of h . $\square$

Theorem 12.3. *Let $h' := \frac{h}{\gcd(d_1, \dots, d_n)}$. The centres of $B(W)$ and W are cyclic, generated respectively by $\delta^{h'}$ and $c^{h'}$.*

Proof. That ZW is cyclic of order $\gcd(d_1, \dots, d_n)$ is classical; any central element is regular; because $c^{h'}$ is regular and has the right order, it must generate ZW .

Let us prove study the conjugacy action of δ . Write a given $b \in B(W)$ in Garside normal form $b = \delta^k s_1 \dots s_l$ where $k \in \mathbb{Z}$ and $s_1, \dots, s_l \in [1, c] - \{1, c\}$. The normal form of b^δ is $\delta^k s_1^c \dots s_l^c$.

Because $c^{h'}$ is the smallest central power of c in W , $\delta^{h'}$ is the smallest central power of δ in $B(W)$. Moreover, if $b \in ZB(W)$, then it must commute with δ and any simple term s_i in its normal form $\delta^k s_1 \dots s_l$ must commute with c . Because c is Coxeter element, the centraliser of c in W is $\langle c \rangle$ (this follows from Theorem 1.9 and, actually, extends Corollary 4.4 in [41]). Combining this with Lemma 12.2, we see that $l = 0$, i.e., that $b \in \langle \delta \rangle$. $\square$

Remark 12.4. This generalises [18, Theorem 2.24] and was conjectured by Broué-Malle-Rouquier. When W is badly-generated, the centres should still be cyclic. With the results of [18], the only case left to check is G_{31} . The Garside category structure provided by our previous section should certainly be applicable to obtain this case, but we haven't checked the details of the computation. It should also not be too difficult to check that $ZP(W) = \langle \tau \rangle$.

Because of Theorem 12.3, an element $\gamma \in B(W)$ is periodic if and only if it is *commensurable* with τ (or δ), i.e., if there exists p, q such that

$$\gamma^q = \tau^p.$$

To simplify notations, we restrict our attention to the situation where

$$\gamma^d = \tau$$

and call such a periodic γ a *d-th root of τ* . The theory works the same way for other (p, q) .

When d is regular and x_0 is a ζ_d -regular eigenvector, we may consider the *standard d-th root of τ* represented by

$$\begin{aligned} [0, 1] &\longrightarrow W \setminus V^{\text{reg}} \\ t &\longmapsto e^{2\pi\sqrt{-1}\frac{t}{d}} x_0 \end{aligned}$$

and denoted by $\sqrt[d]{\tau}$. This of course involves choosing a particular basepoint, but because the statements below are “up to conjugacy”, one should not worry too much about this.

A particular case is

$$\sqrt[h]{\tau} = \delta.$$

Theorem 12.5 (Springer theory in braid groups). *Let d be a positive integer.*

- (i) *There exists d-th roots of τ if and only if d is regular.*
- (ii) *When d is regular, there is a single conjugacy class of d-th roots of τ in $B(W)$. In particular, all d-th roots of τ are conjugate to $\sqrt[d]{\tau}$.*
- (iii) *Let ρ be a d-th root of τ . Let w be the image of ρ in W . Then w is ζ_d -regular, and the centraliser $C_{B(W)}(\rho)$ is isomorphic to the braid group $B(W')$ of the centraliser $W' := C_W(w)$.*

Proof. (i) A consequence of [5, Corollary 10.4] is that, if $\tau = \delta^h$ admits d -th roots, then $M_d^{\phi_d^h}$ is non-empty. By Lemma 11.13, this implies that d is regular. The converse is obvious (we may consider the particular root $\sqrt[d]{\tau}$).

(ii) Using [5, Corollary 10.4] and [5, Proposition 9.8], we see that conjugacy classes of d -th roots of τ are in one-to-one correspondence with connected components of the category G' . By Corollary 11.21, this category is connected.

(iii) That w is regular follows from (ii), because it is conjugate to the image in W of $\sqrt[d]{\tau}$, whose image in W has a x_0 as ζ_d -regular eigenvector. The assertion about the centraliser follows from its categorical rephrasing in M_d , where it is trivial (the conjugacy action being a power of the diagram automorphism of the Garside structure). $\square$

Remark 12.6. This answers many questions and conjectures by Broué, Michel and others (see [17] for more details). Particular cases of (i) were obtained by Broué-Michel and, independently, by Shvartsman, [19, 43]. Assertion (ii) can be viewed either as a Kerékjártó type theorem (“*all periodic elements are conjugate to a rotation*”, see [5]) or as a Sylow type theorem (“*all $C_{B(W)}(\rho)$ are conjugate*”). The type A case of (ii) actually follows from Kerékjártó’s theorem on periodic homeomorphism of the disk. The type A case of (iii) was proved in [7].

Of course, the most natural interpretation is to view the theorem as providing a braid analog of Theorem 1.9.

Remark 12.7. Let W' be the centraliser of a regular element in W . Let W'' be the centraliser of a regular element in W' . In terms of orbit varieties, $W' \backslash V' = (W \backslash V)^{\mu_d}$ and $W'' \backslash V'' = (W \backslash V)^{\mu_{de}}$. Regular elements of W' are regular in W . It probably not difficult, by applying Theorem 12.5 to the pair (W, W'') , to generalise the result to the pair (W', W'') .

13. GENERALISED NON-CROSSING PARTITIONS

Here again, W is an irreducible well-generated complex reflection group.

When W is of type A_{n-1} , the lattice $(S, \preceq)$ is isomorphic to the lattice of non-crossing partitions of a regular n -gon ([7], [12]). Following [37], [3] and [6], we call *lattice of generalised non-crossing partitions of type W* the lattice

$$(S, \preceq),$$

and *Catalan number of type W* the number

$$\text{Cat}(W) := \prod_{i=1}^n \frac{d_i + h}{d_i}.$$

The operation sending $s \preceq t$ to $s^{-1}t$ is an analogue of the Kreweras complement operation. The map $s \mapsto s^{-1}\delta$ is an anti-automorphism of the lattice.

In the Coxeter case, Chapoton (see [20]) discovered a general formula for the number $Z_W(N)$ of weak chains $s_1 \preceq s_2 \preceq \cdots \preceq s_{N-1}$ of length $N - 1$ in $(S, \preceq)$, or equivalently for the number of *complete* weak chains

$$s_1 \preceq s_2 \preceq \cdots \preceq s_{N-1} \preceq s_N = \delta$$

of length N . This formula continues to hold, though we are only able to prove this case-by-case (see [1] and [20] for the Coxeter types; the $G(e, e, n)$ case was done in [6]; the remaining types are done by computer).

Proposition 13.1. *We have, for all N ,*

$$Z_W(N) = \prod_{i=1}^n \frac{d_i + (N - 1)h}{d_i}.$$

Corollary 13.2. *We have $|S| = \text{Cat}(W)$.*

Another interesting numerical invariant is the *Poincaré polynomial*

$$\text{Poin}(S) := \sum_{s \in S} t^{\ell(s)}.$$

W	degrees	$ R $	$\text{Cat}(W)$	$\text{Poin}(S)$	$ \text{Red}_R(c) $
$G_{23} (H_3)$	2, 6, 10	15	32	$1 + 15t + 15t^2 + t^3$	50
G_{24}	4, 6, 14	21	30	$1 + 14t + 14t^2 + t^3$	49
G_{27}	6, 12, 30	45	42	$1 + 20t + 20t^2 + t^3$	75
$G_{28} (F_4)$	2, 6, 8, 12	24	105	$1 + 24t + 55t^2 + 24t^3 + t^4$	432
G_{29}	4, 8, 12, 20	40	112	$1 + 25t + 60t^2 + 25t^3 + t^4$	500
$G_{30} (H_4)$	2, 12, 20, 30	60	280	$1 + 60t + 158t^2 + 60t^3 + t^4$	1350
G_{33}	4, 6, 10, 12, 18	45	308	$1 + 30t + 123t^2 + 123t^3 + 30t^4 + t^5$	4374
G_{34}	6, 12, 18, 24, 30, 42	126	1584	$1 + 56t + 385t^2 + 700t^3 + 385t^4 + 56t^5 + t^6$	100842
$G_{35} (E_6)$	2, 5, 6, 8, 9, 12	36	833	$1 + 36t + 204t^2 + 351t^3 + 204t^4 + 36t^5 + t^6$	41472
$G_{36} (E_7)$	2, 6, 8, 10, 12, 14, 18	63	4160	$1 + 63t + 546t^2 + 1470t^3 + 1470t^4 + 546t^5 + 63t^6 + t^7$	1062882
$G_{37} (E_8)$	2, 8, 12, 14, 18, 20, 24, 30	120	25080	$1 + 120t + 1540t^2 + 6120t^3 + 9518t^4 + 6120t^5 + 1540t^6 + 120t^7 + t^8$	37968750

TABLE 2. Numerical invariants of generalised non-crossing partitions.

The numerical data for the exceptional types (real and non-real) is summarised in Table 2. The coefficient of t in the Poincaré polynomial is the cardinal of R_c . One observes that

$$R = R_c \Leftrightarrow W \text{ is real.}$$

In the Weyl group case, $\text{Poin}(S)$ may be interpreted as the Poincaré polynomial of the cohomology of a toric variety related to cluster algebras ([20]).

When W' is the (not necessarily well-generated) centraliser of a d -regular element in a well-generated W , the natural substitute for $Z_W(N)$ (and for $\text{Cat}(W) = Z_W(2)$) is the number $Z'_{W'}(N)$ of complete weak N -chains of simples in the reduced dual Garside category M' associated with W' (see Definition 11.14). As in Section 11, let $d' := d/d \wedge h$. Complete weak N -chains of simples in the d' -divided category $M_{d'}$ may be identified with complete weak Nd' -chains of simples in M . There is a natural μ_d -action on these chains. The number $Z'_{W'}(N)$ is the number of μ_d -fixed complete weak Nd' -chains of simples in M . In a joint work with Vic Reiner, we conjecture that the μ_d -action exhibits a *cyclic sieving phenomenon*: the number of fixed points should be the value at ζ_d of a q -analog of the number of chains:

Conjecture 13.3 (Conjecture 6.5 in [9]). *Let q be an indeterminate. For any $a \in \mathbb{Z}_{\geq 1}$, set $[a]_q := 1 + q + \dots + q^{a-1}$. Then*

$$\prod_{i=1}^n \frac{[d_i + (Nd' - 1)h]_q}{[d_i]_q}$$

is a polynomial in q whose value at $q = \zeta_d$ is $Z'_{W'}(N)$.

When $N = 1$, the formula should give the number of objects in M' .

Example 13.4. When the complex reflection group $W' = G_{31}$ is viewed as the centraliser of a 4-regular element in a Weyl group of type E_8 , one checks that the number of objects in M' is 88, as predicted by the conjecture.

ACKNOWLEDGEMENTS

Work on this project started during stimulating visits to KIAS (Seoul) and RIMS (Kyoto), in the summer of 2003, when I realised that the dual braid monoid construction extends to duality groups. I thank Sang Jin Lee and Kyoji Saito for their hospitality and for their interests in discussing these subjects. After this initial progress, I remained stucked for many months, trying to construct the open covering using convex geometry in V^{reg} (refining [3, Section 4]). Two observations helped me figure out that working in the quotient space was more appropriate. First, Kyoji Saito pointed out that the starting point for the construction of the *flat structure* (or *Frobenius manifold structure*, see [39]) on real reflection orbifolds was precisely the duality between degrees and codegrees. The intuition that the flat structure has something to do with the $K(\pi, 1)$ property is explicitly mentioned as a motivation for [39] (see also [40, p. 7]). The second useful discussion was with Frédéric Chapoton, who pointed out the numerical coincidence, in the Coxeter case, between the degree of the Lyashko-Looijenga covering and the number of maximal chains in the lattice of non-crossing partitions.

I thank Pierre Deligne, Eduard Looijenga, Jean Michel, Vic Reiner and Vivien Ripoll for comments, critics and suggestions.

Theorem 12.5 is an overdue answer to a question Michel Broué asked me in 1997, when I was a graduate student under his supervision. Most of my work on braid groups was motivated by this problem. I am glad to have been able to solve it just in time for his sixtieth birthday.

REFERENCES

- [1] Ch. Athanasiadis, V. Reiner, *Non-crossing partitions for the group D_n* , SIAM J. Discrete Math. **18** (2004), 397–417.
- [2] D. Bessis, *Zariski theorems and diagrams for braid groups*, Invent. Math. **145** (2001), 487–507.
- [3] D. Bessis, *The dual braid monoid*, Ann. Sci. École Norm. Sup. **36** (2003), 647–683.
- [4] D. Bessis, *A dual braid monoid for the free group*, J. Algebra **302** (2006), 55–69.
- [5] D. Bessis, *Garside categories, periodic loops and cyclic sets*, arXiv:math.GR/0610778.
- [6] D. Bessis, R. Corran, *Non-crossing partitions of type (e, e, r)* , Advances Math. **202** (2006), 1–49.
- [7] D. Bessis, F. Digne, J. Michel, *Springer theory in braid groups and the Birman-Ko-Lee monoid*, Pacific J. Math. **205** (2002), 287–310.
- [8] D. Bessis, J. Michel, *Explicit presentations for exceptional braid groups*, Experiment. Math. **13** (2004), 257–266.
- [9] D. Bessis, V. Reiner, *Cyclic sieving of noncrossing partitions for complex reflection groups*, arXiv:math.CO/0701792.
- [10] M. Bestvina, *Non-positively curved aspects of Artin groups of finite type*, Geometry and Topology **3** (1999), 269–302.
- [11] J. Birman, K. H. Ko, S. J. Lee, *A new approach to the word and conjugacy problem in the braid groups*, Adv. Math. **139** (1998), no. 2, 322–353.
- [12] T. Brady, *A partial order on the symmetric group and new $K(\pi, 1)$ ’s for the braid groups*, Adv. Math. **161** (2001), 20–40.
- [13] T. Brady, C. Watt, *$K(\pi, 1)$ ’s for Artin groups of finite type*, Geom. Dedicata **94** (2002), 225–250.
- [14] T. Brady, C. Watt, *Lattices in finite real reflection groups*, arXiv:math.CO/0501502.
- [15] E. Brieskorn, *Die Fundamentalgruppe des Raumes der regulären Orbits einer endlichen komplexen Spiegelungsgruppe*, Invent. Math. **12** (1971), 57–61.
- [16] E. Brieskorn, *Sur les groupe de tresses*, Séminaire Bourbaki 1971/72, Lectures Notes in Math. **317**, Springer-Verlag, 1973, 21–44.
- [17] M. Broué, *Reflection groups, braid groups, Hecke algebras, finite reductive groups*, Current Developments in Mathematics 2000 (Harvard university and MIT), International Press, Boston (2001), 1–103.
- [18] M. Broué, G. Malle, R. Rouquier, *Complex reflection groups, braid groups, Hecke algebras*, J. reine angew. Math. **500** (1998), 127–190.
- [19] M. Broué, J. Michel, *Sur certains éléments réguliers des groupes de Weyl et les variétés de Deligne-Lusztig associées*, Proceedings de la Semaine de Luminy “Représentations des groupes réductifs finis”, 73–139, Birkhäuser, 1996.

- [20] F. Chapoton, *Enumerative properties of generalized associahedra*, Sémin. Lothar. Combin. **51** (2004), Art. B51b, 16 pp. (electronic).
- [21] R. Charney, J. Meier, K. Whittlesey, *Bestvina's normal form complex and the homology of Garside groups*, Geom. Dedicata **105** (2004), 171–188.
- [22] W. Couwenberg, G. Heckman, E. Looijenga, *Geometric structures on the complement of a projective arrangement*, Pub. Math. IHES **101** (2005), 69–161.
- [23] P. Dehornoy, *Groupes de Garside*, Ann. Sci. École Norm. Sup. (4) **35** (2002) 267–306
- [24] P. Dehornoy, L. Paris, *Gaussian groups and Garside groups, two generalizations of Artin groups*, Proc. of London Math. Soc. **79** (1999), 569–604.
- [25] P. Deligne, *Les immeubles des groupes de tresses généralisés*, Invent. Math. **17** (1972), 273–302.
- [26] P. Deligne, letter to E. Looijenga, 9/3/1974.
- [27] F. Digne, J. Michel, *Endomorphisms of Deligne-Lusztig varieties*, Nagoya Math. J. **183** (2006), 35–103.
- [28] E. Fadell, L. Neuwirth, *Configuration spaces*, Math. Scand. **10** (1962), 111–118.
- [29] W. Fulton, *Intersection theory*, Second edition, Springer-Verlag, 1998.
- [30] A. Hatcher, *Algebraic topology*, Cambridge University Press, 2002.
- [31] G. Lehrer, J. Michel, *Invariant theory and eigenspaces for unitary reflection groups*, C. R. Math. Acad. Sci. Paris **336** (2003), no. 10, 795–800.
- [32] E. Looijenga, *The complement of the bifurcation variety of a simple singularity*, Invent. Math. **23** (1974), 105–116.
- [33] T. Nakamura, *A note of the $K(\pi, 1)$ -property of the orbit space of the unitary reflection group $G(m, l, n)$* , Sci. Papers College of Arts and Sciences, Univ. Tokyo **33** (1983), 1–6.
- [34] P. Orlik, L. Solomon, *Unitary reflection groups and cohomology*, Invent. Math. **59** (1980), 77–94.
- [35] P. Orlik, L. Solomon, *Discriminants in the invariant theory of reflection groups*, Nagoya Math. J. **109** (1988), 23–45.
- [36] P. Orlik, H. Terao, *Arrangements of hyperplanes*, Grundlehren der mathematischen Wissenschaften **300**, Springer-Verlag, 1992.
- [37] V. Reiner, *Non-crossing partitions for classical reflection groups*, Discrete Math. **177** (1997), 195–222
- [38] K. Saito, *On a linear structure of the quotient variety by a finite reflection group*, Publ. Res. Inst. Math. Sci. Kyoto Univ. **29** (1993), 535–579.
- [39] K. Saito, *Uniformization of the orbifold of a finite reflection group*, RIMS preprint, may 2003.
- [40] K. Saito, *The polyhedra dual to the Weyl chamber decomposition: a précis*, RIMS preprint, september 2004.
- [41] T. A. Springer, *Regular elements of finite reflection groups*, Invent. Math. **25** (1974), 159–198.
- [42] G. C. Shephard, J. A. Todd, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274–304.
- [43] O. V. Shvartsman, *Torsion in the quotient group of the Artin-Brieskorn braid group with respect to the center, and regular Springer numbers* (Russian), Funktsional. Anal. i Prilozhen. **30** (1996), 39–46; translation in Funct. Anal. Appl. **30** (1996), no. 1, 30–35.

DMA, ÉCOLE NORMALE SUPÉRIEURE, 45 RUE D'ULM, 75230 PARIS CEDEX 05, FRANCE

E-mail address: david dot bessis at ens dot fr